

A SEMI-SMALL DECOMPOSITION OF THE CHOW RING OF A MATROID

TOM BRADEN, JUNE HUH, JACOB P. MATHERNE, NICHOLAS PROUDFOOT, AND BOTONG WANG

ABSTRACT. We give a semi-small orthogonal decomposition of the Chow ring of a matroid M . The decomposition is used to give simple proofs of Poincaré duality, the hard Lefschetz theorem, and the Hodge–Riemann relations for the Chow ring, recovering the main result of [AHK18]. We also show that a similar semi-small orthogonal decomposition holds for the augmented Chow ring of M .

1. INTRODUCTION

A *matroid* M on a finite set E is a nonempty collection of subsets of E , called *flats* of M , that satisfies the following properties:

- (1) The intersection of any two flats is a flat.
- (2) For any flat F , any element in $E \setminus F$ is contained in exactly one flat that is minimal among the flats strictly containing F .

Throughout, we suppose in addition that M is a *loopless* matroid:

- (3) The empty subset of E is a flat.

We write $\mathcal{L}(M)$ for the lattice of all flats of M . Every maximal flag of proper flats of M has the same cardinality d , called the *rank* of M . A matroid can be equivalently defined in terms of its *independent sets* or the *rank function*. For background in matroid theory, we refer to [Oxl11] and [Wel76].

The first aim of the present paper is to decompose the Chow ring of M as a module over the Chow ring of the deletion $M \setminus i$ (Theorem 1.1). The decomposition resembles the decomposition of the cohomology ring of a projective variety induced by a semi-small map. In Section 4, we use the decomposition to give simple proofs of Poincaré duality, the hard Lefschetz theorem, and the Hodge–Riemann relations for the Chow ring, recovering the main result of [AHK18].

June Huh received support from NSF Grant DMS-1638352 and the Ellentuck Fund. Jacob Matherne received support from NSF Grant DMS-1638352, the Association of Members of the Institute for Advanced Study, and the Max Planck Institute for Mathematics in Bonn. Nicholas Proudfoot received support from NSF Grant DMS-1565036. Botong Wang received support from NSF Grant DMS-1701305 and the Alfred P. Sloan foundation.

The second aim of the present paper is to introduce the augmented Chow ring of M , which contains the graded Möbius algebra of M as a subalgebra. We give an analogous semi-small decomposition of the augmented Chow ring of M as a module over the augmented Chow ring of the deletion $M \setminus i$ (Theorem 1.2), and use this to prove Poincaré duality, the hard Lefschetz theorem, and the Hodge–Riemann relations for the augmented Chow ring. These results will play a major role in the forthcoming paper [BHM⁺], where we will prove the Top-Heavy conjecture along with the nonnegativity of the coefficients of the Kazhdan–Lusztig polynomial of a matroid.

1.1. Let \underline{S}_M be the ring of polynomials with variables labeled by the nonempty proper flats of M :

$$\underline{S}_M := \mathbb{Q}[x_F \mid F \text{ is a nonempty proper flat of } M].$$

The *Chow ring* of M , introduced by Feichtner and Yuzvinsky in [FY04], is the quotient algebra¹

$$\underline{\text{CH}}(M) := \underline{S}_M / (\underline{I}_M + \underline{J}_M),$$

where \underline{I}_M is the ideal generated by the linear forms

$$\sum_{i_1 \in F} x_F - \sum_{i_2 \in F} x_F, \quad \text{for every pair of distinct elements } i_1 \text{ and } i_2 \text{ of } E,$$

and \underline{J}_M is the ideal generated by the quadratic monomials

$$x_{F_1} x_{F_2}, \quad \text{for every pair of incomparable nonempty proper flats } F_1 \text{ and } F_2 \text{ of } M.$$

When E is nonempty, the Chow ring of M admits a *degree map*

$$\underline{\text{deg}}_M : \underline{\text{CH}}^{d-1}(M) \longrightarrow \mathbb{Q}, \quad x_{\mathcal{F}} := \prod_{F \in \mathcal{F}} x_F \longmapsto 1,$$

where \mathcal{F} is any complete flag of nonempty proper flats of M (Definition 2.12). For any integer k , the degree map defines the *Poincaré pairing*

$$\underline{\text{CH}}^k(M) \times \underline{\text{CH}}^{d-k-1}(M) \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto \underline{\text{deg}}_M(\eta_1 \eta_2).$$

If M is representable over a field,² then the Chow ring of M is isomorphic to the Chow ring of a smooth projective variety over the field (Remark 2.13).

Let i be an element of E , and let $M \setminus i$ be the *deletion* of i from M . By definition, $M \setminus i$ is the matroid on $E \setminus i$ whose flats are the sets of the form $F \setminus i$ for a flat F of M . The Chow rings of M and $M \setminus i$ are related by the graded algebra homomorphism

$$\underline{\theta}_i = \underline{\theta}_i^M : \underline{\text{CH}}(M \setminus i) \longrightarrow \underline{\text{CH}}(M), \quad x_F \longmapsto x_F + x_{F \cup i},$$

¹A slightly different presentation for the Chow ring of M was used in [FY04] in a more general context. The present description was used in [AHK18], where the Chow ring of M was denoted $A(M)$. For a comparison of the two presentations, see [BES].

²We say that M is *representable* over a field \mathbb{F} if there exists a linear subspace $V \subseteq \mathbb{F}^E$ such that $S \subseteq E$ is independent if and only if the projection from V to \mathbb{F}^S is surjective. Almost all matroids are not representable over any field [Nel18].

where a variable in the target is set to zero if its label is not a flat of M . Let $\underline{\text{CH}}_{(i)}$ be the image of the homomorphism θ_i , and let $\underline{\mathcal{S}}_i$ be the collection

$$\underline{\mathcal{S}}_i = \underline{\mathcal{S}}_i(M) = \{F \mid F \text{ is a nonempty proper subset of } E \setminus i \text{ such that } F \in \mathcal{L}(M) \text{ and } F \cup i \in \mathcal{L}(M)\}.$$

The element i is said to be a *coloop* of M if the ranks of M and $M \setminus i$ are not equal.

Theorem 1.1. *If i is not a coloop of M , there is a direct sum decomposition of $\underline{\text{CH}}(M)$ into indecomposable graded $\underline{\text{CH}}(M \setminus i)$ -modules*

$$\underline{\text{CH}}(M) = \underline{\text{CH}}_{(i)} \oplus \bigoplus_{F \in \underline{\mathcal{S}}_i} x_{F \cup i} \underline{\text{CH}}_{(i)}. \quad (\underline{\text{D}}_1)$$

All pairs of distinct summands are orthogonal for the Poincaré pairing of $\underline{\text{CH}}(M)$. If i is a coloop of M , there is a direct sum decomposition of $\underline{\text{CH}}(M)$ into indecomposable graded $\underline{\text{CH}}(M \setminus i)$ -modules³

$$\underline{\text{CH}}(M) = \underline{\text{CH}}_{(i)} \oplus x_{E \setminus i} \underline{\text{CH}}_{(i)} \oplus \bigoplus_{F \in \underline{\mathcal{S}}_i} x_{F \cup i} \underline{\text{CH}}_{(i)}. \quad (\underline{\text{D}}_2)$$

All pairs of distinct summands except for the first two are orthogonal for the Poincaré pairing of $\underline{\text{CH}}(M)$.

We write $\text{rk}_M : 2^E \rightarrow \mathbb{N}$ for the rank function of M . For any proper flat F of M , we set⁴

$M^F :=$ the localization of M at F , a loopless matroid on F of rank equal to $\text{rk}_M(F)$,

$M_F :=$ the contraction of M by F , a loopless matroid on $E \setminus F$ of rank equal to $d - \text{rk}_M(F)$.

The lattice of flats of M^F can be identified with the lattice of flats of M that are contained in F , and the lattice of flats of M_F can be identified with the lattice of flats of M that contain F . The $\underline{\text{CH}}(M \setminus i)$ -module summands in the decompositions $(\underline{\text{D}}_1)$ and $(\underline{\text{D}}_2)$ admit isomorphisms

$$\underline{\text{CH}}_{(i)} \cong \underline{\text{CH}}(M \setminus i) \text{ and } x_{F \cup i} \underline{\text{CH}}_{(i)} \cong \underline{\text{CH}}(M_{F \cup i}) \otimes \underline{\text{CH}}(M^F)[-1],$$

where $[-1]$ indicates a degree shift (Propositions 3.4 and 3.5). In addition, if i is a coloop of M ,

$$x_{E \setminus i} \underline{\text{CH}}_{(i)} \cong \underline{\text{CH}}(M \setminus i)[-1].$$

Numerically, the semi-smallness of the decomposition $(\underline{\text{D}}_1)$ is reflected in the identity

$$\dim x_{F \cup i} \underline{\text{CH}}_{(i)}^{k-1} = \dim x_{F \cup i} \underline{\text{CH}}_{(i)}^{d-k-2} \text{ for } F \in \underline{\mathcal{S}}_i.$$

³When $E = \{i\}$, we treat the symbol x_\emptyset as zero in the right-hand side of $(\underline{\text{D}}_2)$.

⁴The symbols M^F and M_F appear inconsistently in the literature, sometimes this way and sometimes interchanged. The localization is frequently called the restriction. On the other hand, the contraction is also sometimes called the restriction, especially in the context of hyperplane arrangements, so we avoid the word restriction to minimize ambiguity.

When M is the Boolean matroid on E , the graded dimension of $\underline{\text{CH}}(M)$ is given by the Eulerian numbers $\langle \binom{d}{k} \rangle$, and the decomposition $(\underline{\text{D}}_2)$ specializes to the known quadratic recurrence relation

$$s_d(t) = s_{d-1}(t) + t \sum_{k=0}^{d-2} \binom{d-1}{k} s_k(t) s_{d-k-1}(t), \quad s_0(t) = 1,$$

where $s_k(t)$ is the k -th Eulerian polynomial [Pet15, Theorem 1.5].

1.2. We also give similar decompositions for the augmented Chow ring of M , which we now introduce. Let S_M be the ring of polynomials in two sets of variables

$$S_M := \mathbb{Q}[y_i \mid i \text{ is an element of } E] \otimes \mathbb{Q}[x_F \mid F \text{ is a proper flat of } M].$$

The *augmented Chow ring* of M is the quotient algebra

$$\text{CH}(M) := S_M / (I_M + J_M),$$

where I_M is the ideal generated by the linear forms

$$y_i - \sum_{i \notin F} x_F, \quad \text{for every element } i \text{ of } E,$$

and J_M is the ideal generated by the quadratic monomials

$$\begin{aligned} & x_{F_1} x_{F_2}, \quad \text{for every pair of incomparable proper flats } F_1 \text{ and } F_2 \text{ of } M, \text{ and} \\ & y_i x_F, \quad \text{for every element } i \text{ of } E \text{ and every proper flat } F \text{ of } M \text{ not containing } i. \end{aligned}$$

The augmented Chow ring of M admits a *degree map*

$$\deg_M : \text{CH}^d(M) \longrightarrow \mathbb{Q}, \quad x_{\mathcal{F}} := \prod_{F \in \mathcal{F}} x_F \longmapsto 1,$$

where \mathcal{F} is any complete flag of proper flats of M (Definition 2.12). For any integer k , the degree map defines the *Poincaré pairing*

$$\text{CH}^k(M) \times \text{CH}^{d-k}(M) \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto \deg_M(\eta_1 \eta_2).$$

If M is representable over a field, then the augmented Chow ring of M is isomorphic to the Chow ring of a smooth projective variety over the field (Remark 2.13). The augmented Chow ring contains the *graded Möbius algebra* $H(M)$ (Proposition 2.15), and it is related to the Chow ring of M by the isomorphism

$$\underline{\text{CH}}(M) \cong \text{CH}(M) \otimes_{H(M)} \mathbb{Q}.$$

The $H(M)$ -module structure of $\text{CH}(M)$ will be studied in detail in the forthcoming paper [BHM⁺].

As before, we write $M \setminus i$ for the matroid obtained from M by deleting the element i . The augmented Chow rings of M and $M \setminus i$ are related by the graded algebra homomorphism

$$\theta_i = \theta_i^M : \text{CH}(M \setminus i) \longrightarrow \text{CH}(M), \quad x_F \longmapsto x_F + x_{F \cup i},$$

where a variable in the target is set to zero if its label is not a flat of M . Let $\text{CH}_{(i)}$ be the image of the homomorphism θ_i , and let \mathcal{S}_i be the collection

$$\mathcal{S}_i = \mathcal{S}_i(M) := \{F \mid F \text{ is a proper subset of } E \setminus i \text{ such that } F \in \mathcal{L}(M) \text{ and } F \cup i \in \mathcal{L}(M)\}.$$

Theorem 1.2. If i is not a coloop of M , there is a direct sum decomposition of $\text{CH}(M)$ into indecomposable graded $\text{CH}(M \setminus i)$ -modules

$$\text{CH}(M) = \text{CH}_{(i)} \oplus \bigoplus_{F \in \mathcal{S}_i} x_{F \cup i} \text{CH}_{(i)}. \quad (\text{D}_1)$$

All pairs of distinct summands are orthogonal for the Poincaré pairing of $\text{CH}(M)$. If i is a coloop of M , there is a direct sum decomposition of $\text{CH}(M)$ into indecomposable graded $\text{CH}(M \setminus i)$ -modules

$$\text{CH}(M) = \text{CH}_{(i)} \oplus x_{E \setminus i} \text{CH}_{(i)} \oplus \bigoplus_{F \in \mathcal{S}_i} x_{F \cup i} \text{CH}_{(i)}. \quad (\text{D}_2)$$

All pairs of distinct summands except for the first two are orthogonal for the Poincaré pairing of $\text{CH}(M)$.

The $\text{CH}(M \setminus i)$ -module summands in the decompositions (D_1) and (D_2) admit isomorphisms

$$\text{CH}_{(i)} \cong \text{CH}(M \setminus i) \quad \text{and} \quad x_{F \cup i} \text{CH}_{(i)} \cong \underline{\text{CH}}(M_{F \cup i}) \otimes \text{CH}(M^F)[-1],$$

where $[-1]$ indicates a degree shift (Propositions 3.4 and 3.5). In addition, if i is a coloop of M ,

$$x_{E \setminus i} \text{CH}_{(i)} \cong \text{CH}(M \setminus i)[-1].$$

Numerically, the semi-smallness of the decomposition (D_1) is reflected in the identity

$$\dim x_{F \cup i} \text{CH}_{(i)}^{k-1} = \dim x_{F \cup i} \text{CH}_{(i)}^{d-k-1} \quad \text{for } F \in \mathcal{S}_i.$$

1.3. Let B be the Boolean matroid on E . By definition, every subset of E is a flat of B . The Chow rings of B and M are related by the surjective graded algebra homomorphism

$$\underline{\text{CH}}(B) \longrightarrow \underline{\text{CH}}(M), \quad x_S \longmapsto x_S,$$

where a variable in the target is set to zero if its label is not a flat of M . Similarly, we have a surjective graded algebra homomorphism

$$\text{CH}(B) \longrightarrow \text{CH}(M), \quad x_S \longmapsto x_S,$$

where a variable in the target is set to zero if its label is not a flat of M . As in [AHK18, Section 4], we may identify the Chow ring $\underline{\text{CH}}(B)$ with the ring of piecewise polynomial functions modulo linear functions on the normal fan $\underline{\Pi}_B$ of the standard permutohedron in \mathbb{R}^E . Similarly, the augmented Chow ring $\text{CH}(B)$ can be identified with the ring of piecewise polynomial functions modulo linear functions of the normal fan Π_B of the stellahedron in \mathbb{R}^E (Definition 2.4). A convex piecewise

linear function on a complete fan is said to be *strictly convex* if there is a bijection between the cones in the fan and the faces of the graph of the function.

In Section 4, we use Theorems 1.1 and 1.2 to give simple proofs of Poincaré duality, the hard Lefschetz theorem, and the Hodge–Riemann relations for $\underline{\text{CH}}(\text{M})$ and $\text{CH}(\text{M})$.

Theorem 1.3. Let $\underline{\ell}$ be a strictly convex piecewise linear function on Π_{B} , viewed as an element of $\underline{\text{CH}}^1(\text{M})$.

(1) (Poincaré duality theorem) For every nonnegative integer $k < \frac{d}{2}$, the bilinear pairing

$$\underline{\text{CH}}^k(\text{M}) \times \underline{\text{CH}}^{d-k-1}(\text{M}) \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto \underline{\text{deg}}_{\text{M}}(\eta_1 \eta_2)$$

is non-degenerate.

(2) (Hard Lefschetz theorem) For every nonnegative integer $k < \frac{d}{2}$, the multiplication map

$$\underline{\text{CH}}^k(\text{M}) \longrightarrow \underline{\text{CH}}^{d-k-1}(\text{M}), \quad \eta \longmapsto \underline{\ell}^{d-2k-1} \eta$$

is an isomorphism.

(3) (Hodge–Riemann relations) For every nonnegative integer $k < \frac{d}{2}$, the bilinear form

$$\underline{\text{CH}}^k(\text{M}) \times \underline{\text{CH}}^k(\text{M}) \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto (-1)^k \underline{\text{deg}}_{\text{M}}(\underline{\ell}^{d-2k-1} \eta_1 \eta_2)$$

is positive definite on the kernel of multiplication by $\underline{\ell}^{d-2k}$.

Let ℓ be a strictly convex piecewise linear function on Π_{B} , viewed as an element of $\text{CH}^1(\text{M})$.

(4) (Poincaré duality theorem) For every nonnegative integer $k \leq \frac{d}{2}$, the bilinear pairing

$$\text{CH}^k(\text{M}) \times \text{CH}^{d-k}(\text{M}) \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto \text{deg}_{\text{M}}(\eta_1 \eta_2)$$

is non-degenerate.

(5) (Hard Lefschetz theorem) For every nonnegative integer $k \leq \frac{d}{2}$, the multiplication map

$$\text{CH}^k(\text{M}) \longrightarrow \text{CH}^{d-k}(\text{M}), \quad \eta \longmapsto \ell^{d-2k} \eta$$

is an isomorphism.

(6) (Hodge–Riemann relations) For every nonnegative integer $k \leq \frac{d}{2}$, the bilinear form

$$\text{CH}^k(\text{M}) \times \text{CH}^k(\text{M}) \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto (-1)^k \text{deg}_{\text{M}}(\ell^{d-2k} \eta_1 \eta_2)$$

is positive definite on the kernel of multiplication by ℓ^{d-2k+1} .

Theorem 1.3 holds non-vacuously, as there are strictly convex piecewise linear functions on Π_B and $\underline{\Pi}_B$ (Proposition 2.6). The first part of Theorem 1.3 on $\underline{\text{CH}}(M)$ recovers the main result of [AHK18].⁵ The second part of Theorem 1.3 on $\text{CH}(M)$ is new.

1.4. In Section 5, we use Theorems 1.1 and 1.2 to obtain decompositions of $\underline{\text{CH}}(M)$ and $\text{CH}(M)$ related to those appearing in [AHK18, Theorem 6.18]. Let $\underline{H}_\alpha(M)$ be the subalgebra of $\underline{\text{CH}}(M)$ generated by the element

$$\underline{\alpha}_M := \sum_{i \in G} x_G \in \underline{\text{CH}}^1(M),$$

where the sum is over all nonempty proper flats G of M containing a given element i in E , and let $H_\alpha(M)$ be the subalgebra of $\text{CH}(M)$ generated by the element

$$\alpha_M := \sum_G x_G \in \text{CH}^1(M),$$

where the sum is over all proper flats G of M . We define graded subspaces $\underline{J}_\alpha(M)$ and $J_\alpha(M)$ by

$$\underline{J}_\alpha^k(M) := \begin{cases} \underline{H}_\alpha^k(M) & \text{if } k \neq d-1, \\ 0 & \text{if } k = d-1, \end{cases} \quad J_\alpha^k(M) := \begin{cases} H_\alpha^k(M) & \text{if } k \neq d, \\ 0 & \text{if } k = d. \end{cases}$$

A degree computation shows that the elements $\underline{\alpha}_M^{d-1}$ and α_M^d are nonzero (Proposition 2.26).

Theorem 1.4. Let $\mathcal{C} = \mathcal{C}(M)$ be the set of all nonempty proper flats of M , and let $\underline{\mathcal{C}} = \underline{\mathcal{C}}(M)$ be the set of all proper flats of M with rank at least two.

(1) We have a decomposition of $\underline{H}_\alpha(M)$ -modules

$$\underline{\text{CH}}(M) = \underline{H}_\alpha(M) \oplus \bigoplus_{F \in \underline{\mathcal{C}}} \psi_M^F \underline{\text{CH}}(M_F) \otimes \underline{J}_\alpha(M^F). \quad (\text{D}_3)$$

All pairs of distinct summands are orthogonal for the Poincaré pairing of $\underline{\text{CH}}(M)$.

(2) We have a decomposition of $H_\alpha(M)$ -modules

$$\text{CH}(M) = H_\alpha(M) \oplus \bigoplus_{F \in \mathcal{C}} \psi_M^F \text{CH}(M_F) \otimes J_\alpha(M^F). \quad (\text{D}_3)$$

All pairs of distinct summands are orthogonal for the Poincaré pairing of $\text{CH}(M)$.

Here ψ_M^F is the injective $\underline{\text{CH}}(M)$ -module homomorphism (Propositions 2.21 and 2.22)

$$\psi_M^F : \underline{\text{CH}}(M_F) \otimes \underline{\text{CH}}(M^F) \longrightarrow \underline{\text{CH}}(M), \quad \prod_{F'} x_{F' \setminus F} \otimes \prod_{F''} x_{F''} \longmapsto x_F \prod_{F'} x_{F'} \prod_{F''} x_{F''},$$

⁵Independent proofs of Poincaré duality for $\underline{\text{CH}}(M)$ were given in [BES] and [BDF]. The authors of [BES] also prove the degree 1 Hodge–Riemann relations for $\underline{\text{CH}}(M)$.

and ψ_M^F is the injective $\text{CH}(M)$ -module homomorphism (Propositions 2.18 and 2.19)

$$\psi_M^F : \underline{\text{CH}}(M_F) \otimes \text{CH}(M^{F'}) \longrightarrow \text{CH}(M) \quad \prod_{F'} x_{F' \setminus F} \otimes \prod_{F''} x_{F''} \longmapsto x_F \prod_{F'} x_{F'} \prod_{F''} x_{F''}.$$

When M is the Boolean matroid on E , the decomposition (D₃) specializes to a linear recurrence relation for the Eulerian polynomials

$$0 = 1 + \sum_{k=0}^d \binom{d}{k} \frac{t - t^{d-k}}{1-t} s_k(t), \quad s_0(t) = 1.$$

When applied repeatedly, Theorem 1.4 produces bases of $\underline{\text{CH}}(M)$ and $\text{CH}(M)$ that are permuted by the automorphism group of M .⁶

Acknowledgments. We thank Christopher Eur and Matthew Stevens for useful discussions.

2. THE CHOW RING AND THE AUGMENTED CHOW RING OF A MATROID

In this section, we collect the various properties of the algebras $\underline{\text{CH}}(M)$ and $\text{CH}(M)$ that we will need in order to prove Theorems 1.1–1.4. In Section 2.1, we review the definition and basic properties of the *Bergman fan* and introduce the closely related *augmented Bergman fan* of a matroid. Section 2.2 is devoted to understanding the stars of the various rays in these two fans, while Section 2.3 is where we compute the space of balanced top-dimensional weights on each fan. Feichtner and Yuzvinsky showed that the Chow ring of a matroid coincides with the Chow ring of the toric variety associated with its Bergman fan [FY04, Theorem 3], and we establish the analogous result for the augmented Chow ring in Section 2.4. Section 2.5 is where we show that the augmented Chow ring contains the graded Möbius algebra. In Section 2.6, we use the results of Section 2.2 to construct various homomorphisms that relate the Chow and augmented Chow rings of different matroids.

Remark 2.1. It is worth noting why we need to interpret $\underline{\text{CH}}(M)$ and $\text{CH}(M)$ as Chow rings of toric varieties. First, the study of balanced weights on the Bergman fan and augmented Bergman fan allow us to show that $\underline{\text{CH}}^{d-1}(M)$ and $\text{CH}^d(M)$ are nonzero, which is not easy to prove directly from the definitions. The definition of the pullback and pushforward maps in Section 2.6 is made cleaner by thinking about fans, though it would also be possible to define these maps by taking Propositions 2.17, 2.18, 2.20, 2.21, 2.23, and 2.24 as definitions. Finally, and most importantly, the fan perspective will be essential for understanding the ample classes that appear in Theorem 1.3.

⁶Different bases of $\underline{\text{CH}}(M)$ are given in [FY04, Corollary 1] and [BES, Corollary 3.3.3].

2.1. Fans. Let E be a finite set, and let M be a loopless matroid of rank d on the ground set E . We write rk_M for the rank function of M , and write cl_M for the closure operator of M , which for a set S returns the smallest flat containing S . The *independence complex* I_M of M is the simplicial complex of independent sets of M . A set $I \subseteq E$ is independent if and only if the rank of $\text{cl}_M(I)$ is $|I|$. The vertices of I_M are the elements of the ground set E , and a collection of vertices is a face of I_M when the corresponding set of elements is an independent set of M . The *Bergman complex* $\underline{\Delta}_M$ of M is the order complex of the poset of nonempty proper flats of M . The vertices of $\underline{\Delta}_M$ are the nonempty proper flats of M , and a collection of vertices is a face of $\underline{\Delta}_M$ when the corresponding set of flats is a flag. The independence complex of M is pure of dimension $d - 1$, and the Bergman complex of M is pure of dimension $d - 2$. For a detailed study of the simplicial complexes I_M and $\underline{\Delta}_M$, we refer to [Bjö92]. We introduce the *augmented Bergman complex* Δ_M of M as a simplicial complex that interpolates between the independence complex and the Bergman complex of M .

Definition 2.2. Let I be an independent set of M , and let \mathcal{F} be a flag of proper flats of M . When I is contained in every flat in \mathcal{F} , we say that I is *compatible* with \mathcal{F} and write $I \leq \mathcal{F}$. The *augmented Bergman complex* Δ_M of M is the simplicial complex of all compatible pairs $I \leq \mathcal{F}$, where I is an independent set of M and \mathcal{F} is a flag of proper flats of M .

A vertex of the augmented Bergman complex Δ_M is either a singleton subset of E or a proper flat of M . More precisely, the vertices of Δ_M are the compatible pairs either of the form $\{i\} \leq \emptyset$ or of the form $\emptyset \leq \{F\}$, where i is an element of E and F is a proper flat of M . The augmented Bergman complex contains both the independence complex I_M and the Bergman complex $\underline{\Delta}_M$ as subcomplexes. In fact, Δ_M contains the order complex of the poset of proper flats of M , which is the cone over the Bergman complex with the cone point corresponding to the empty flat. It is straightforward to check that Δ_M is pure of dimension $d - 1$.

Proposition 2.3. The Bergman complex and the augmented Bergman complex of M are both connected in codimension 1.

Proof. The statement about the Bergman complex is a direct consequence of its shellability [Bjö92]. We prove the statement about the augmented Bergman complex using the statement about the Bergman complex.

The claim is that, given any two facets of Δ_M , one may travel from one facet to the other by passing through faces of codimension at most 1. Since the Bergman complex of M is connected in codimension 1, the subcomplex of Δ_M consisting of faces of the form $\emptyset \leq \mathcal{F}$ is connected in codimension 1. Thus it suffices to show that any facet of Δ_M can be connected to a facet of the form $\emptyset \leq \mathcal{F}$ through codimension 1 faces.

Let $I \leq \mathcal{F}$ be a facet of Δ_M . If I is nonempty, choose any element i of I , and consider the flag of flats \mathcal{G} obtained by adjoining the closure of $I \setminus i$ to \mathcal{F} . The independent set $I \setminus i$ is compatible with

the flag \mathcal{G} , and the facet $I \leq \mathcal{F}$ is adjacent to the facet $I \setminus i \leq \mathcal{G}$. Repeating the procedure, we can connect the given facet to a facet of the desired form through codimension 1 faces. \square

Let \mathbb{R}^E be the vector space spanned by the standard basis vectors \mathbf{e}_i corresponding to the elements $i \in E$. For an arbitrary subset $S \subseteq E$, we set

$$\mathbf{e}_S := \sum_{i \in S} \mathbf{e}_i.$$

For an element $i \in E$, we write ρ_i for the ray generated by the vector \mathbf{e}_i in \mathbb{R}^E . For a subset $S \subseteq E$, we write ρ_S for the ray generated by the vector $-\mathbf{e}_{E \setminus S}$ in \mathbb{R}^E , and write $\underline{\rho}_S$ for the ray generated by the vector \mathbf{e}_S in $\mathbb{R}^E / \langle \mathbf{e}_S \rangle$. Using these rays, we construct fan models of the Bergman complex and the augmented Bergman complex as follows.

Definition 2.4. The *Bergman fan* $\underline{\Pi}_M$ of M is a simplicial fan in the quotient space $\mathbb{R}^E / \langle \mathbf{e}_E \rangle$ with rays $\underline{\rho}_F$ for nonempty proper flats F of M . The cones of $\underline{\Pi}_M$ are of the form

$$\underline{\sigma}_{\mathcal{F}} := \text{cone}\{\mathbf{e}_F\}_{F \in \mathcal{F}} = \text{cone}\{-\mathbf{e}_{E \setminus F}\}_{F \in \mathcal{F}},$$

where \mathcal{F} is a flag of nonempty proper flats of M .

The *augmented Bergman fan* Π_M of M is a simplicial fan in \mathbb{R}^E with rays ρ_i for elements i in E and ρ_F for proper flats F of M . The cones of the augmented Bergman fan are of the form

$$\sigma_{I \leq \mathcal{F}} := \text{cone}\{\mathbf{e}_i\}_{i \in I} + \text{cone}\{-\mathbf{e}_{E \setminus F}\}_{F \in \mathcal{F}},$$

where \mathcal{F} is a flag of proper flats of M and I is an independent set of M compatible with \mathcal{F} . We write σ_I for the cone $\sigma_{I \leq \mathcal{F}}$ when \mathcal{F} is the empty flag of flats of M .

Remark 2.5. If E is nonempty, then the Bergman fan $\underline{\Pi}_M$ is the star of the ray ρ_\emptyset in the augmented Bergman fan Π_M . If E is empty, then $\underline{\Pi}_M$ and Π_M both consist of a single 0-dimensional cone.

Let N be another loopless matroid on E . The matroid M is said to be a *quotient* of N if every flat of M is a flat of N . The condition implies that every independent set of M is an independent set of N [Kun86, Proposition 8.1.6]. Therefore, when M is a quotient of N , the augmented Bergman fan of M is a subfan of the augmented Bergman fan of N , and the Bergman fan of M is a subfan of the Bergman fan of N . In particular, we have inclusions of fans $\Pi_M \subseteq \Pi_N$ and $\underline{\Pi}_M \subseteq \underline{\Pi}_N$, where N is the *Boolean matroid* on E defined by the condition that E is an independent set of N .

Proposition 2.6. The Bergman fan and the augmented Bergman fan of N are each normal fans of convex polytopes. In particular, there are strictly convex piecewise linear functions on $\underline{\Pi}_N$ and Π_N .

The above proposition can be used to show that the augmented Bergman fan and the Bergman fan of M are, in fact, fans.




FIGURE 1. The augmented Bergman fan of the rank 2 Boolean matroid on $\{1, 2\}$.

Proof. The statement for the Bergman fan is well-known: The Bergman fan of B is the normal fan of the standard permutohedron in $e_E^\perp \subseteq \mathbb{R}^E$. See, for example, [AHK18, Section 2]. The statement for the augmented Bergman fan Π_B follows from the fact that it is an iterated stellar subdivision of the normal fan of the simplex

$$\text{conv}\{\mathbf{e}_i, \mathbf{e}_E\}_{i \in E} \subseteq \mathbb{R}^E.$$

More precisely, Π_B is isomorphic to the fan $\Sigma_{\mathcal{P}}$ in [AHK18, Definition 2.3], where \mathcal{P} is the order filter of all subsets of $E \cup 0$ containing the new element 0, via the linear isomorphism

$$\mathbb{R}^E \longrightarrow \mathbb{R}^{E \cup 0} / \langle \mathbf{e}_E + \mathbf{e}_0 \rangle, \quad \mathbf{e}_j \longmapsto \mathbf{e}_j.$$

It is shown in [AHK18, Proposition 2.4] that $\Sigma_{\mathcal{P}}$ is an iterated stellar subdivision of the normal fan of the simplex.⁷ \square

A direct inspection shows that Π_M is a *unimodular fan*; that is, the set of primitive ray generators in any cone in Π_M is a subset of a basis of the free abelian group \mathbb{Z}^E . It follows that Π_M is also a unimodular fan; that is, the set of primitive ray generators in any cone in Π_M is a subset of a basis of the free abelian group $\mathbb{Z}^E / \langle \mathbf{e}_E \rangle$.

⁷In fact, the augmented Bergman fan Π_B is the normal fan of the stellahedron in \mathbb{R}^E , the graph associahedron of the star graph with $|E|$ endpoints. We refer to [CD06] and [Dev09] for detailed discussions of graph associahedra and their realizations.

2.2. Stars. For any element i of E , we write $\text{cl}(i)$ for the closure of i in M , and write ι_i for the injective linear map

$$\iota_i : \mathbb{R}^{E \setminus \text{cl}(i)} \longrightarrow \mathbb{R}^E / \langle \mathbf{e}_i \rangle, \quad \mathbf{e}_j \longmapsto \mathbf{e}_j.$$

For any proper flat F of M , we write ι_F for the linear isomorphism

$$\iota_F : \mathbb{R}^{E \setminus F} / \langle \mathbf{e}_{E \setminus F} \rangle \oplus \mathbb{R}^F \longrightarrow \mathbb{R}^E / \langle \mathbf{e}_{E \setminus F} \rangle, \quad \mathbf{e}_j \longmapsto \mathbf{e}_j.$$

For any nonempty proper flat F of M , we write ι_F for the linear isomorphism

$$\iota_F : \mathbb{R}^{E \setminus F} / \langle \mathbf{e}_{E \setminus F} \rangle \oplus \mathbb{R}^F / \langle \mathbf{e}_F \rangle \longrightarrow \mathbb{R}^E / \langle \mathbf{e}_E, \mathbf{e}_{E \setminus F} \rangle, \quad \mathbf{e}_j \longmapsto \mathbf{e}_j.$$

Let M^F be the localization of M at F , and let M_F be the contraction of M by F .

Proposition 2.7. The following are descriptions of the stars of the rays in Π_M and $\underline{\Pi}_M$ using the three linear maps above.

- (1) For any element $i \in E$, the linear map ι_i identifies the augmented Bergman fan of $M_{\text{cl}(i)}$ with the star of the ray ρ_i in the augmented Bergman fan of M :

$$\Pi_{M_{\text{cl}(i)}} \cong \text{star}_{\rho_i} \Pi_M.$$

- (2) For any proper flat F of M , the linear map ι_F identifies the product of the Bergman fan of M_F and the augmented Bergman fan of M^F with the star of the ray ρ_F in the augmented Bergman fan of M :

$$\underline{\Pi}_{M_F} \times \Pi_{M^F} \cong \text{star}_{\rho_F} \Pi_M.$$

- (3) For any nonempty proper flat F of M , the linear map ι_F identifies the product of the Bergman fan of M_F and the Bergman fan of M^F with the star of the ray ρ_F in the Bergman fan of M :

$$\underline{\Pi}_{M_F} \times \underline{\Pi}_{M^F} \cong \text{star}_{\rho_F} \underline{\Pi}_M.$$

Repeated applications of the first statement show that, for any independent set I of M , the star of the cone σ_I in Π_M can be identified with the augmented Bergman fan of $M_{\text{cl}(I)}$, where $\text{cl}(I)$ is the closure of I in M .

Proof. The first statement follows from the following facts: A flat of M contains i if and only if it contains $\text{cl}(i)$, and an independent set of M containing i does not contain any other element in $\text{cl}(i)$. The second and third statements follow directly from the definitions. \square

2.3. Weights. For any simplicial fan Σ , we write Σ_k for the set of k -dimensional cones in Σ . If τ is a codimension 1 face of a cone σ , we write

$\mathbf{e}_{\sigma/\tau} :=$ the primitive generator of the unique ray in σ that is not in τ .

A k -dimensional balanced weight on Σ is a \mathbb{Q} -valued function ω on Σ_k that satisfies the *balancing condition*: For every $(k - 1)$ -dimensional cone τ in Σ ,

$$\sum_{\tau \subset \sigma} \omega(\sigma) \mathbf{e}_{\sigma/\tau} \text{ is contained in the subspace spanned by } \tau,$$

where the sum is over all k -dimensional cones σ containing τ . We write $\text{MW}_k(\Sigma)$ for the group of k -dimensional balanced weights on Σ .

Proposition 2.8. The Bergman fan and the augmented Bergman fan of M have the following unique balancing property.

- (1) A $(d - 1)$ -dimensional weight on Π_M is balanced if and only if it is constant.
- (2) A d -dimensional weight on Π_M is balanced if and only if it is constant.

Proof. The first statement is [AHK18, Proposition 5.2]. We prove the second statement.

Let $\sigma_{I \leq \mathcal{F}}$ be a codimension 1 cone of Π_M , and let F be the smallest flat in $\mathcal{F} \cup \{E\}$. We analyze the primitive generators of the rays in the star of the cone $\sigma_{I \leq \mathcal{F}}$ in Π_M . Let $\text{cl}(I)$ be the closure of I in M . There are two cases.

When the closure of I is not F , the primitive ray generators in question are $-\mathbf{e}_{E \setminus \text{cl}(I)}$ and \mathbf{e}_i , for elements i in F not in the closure of I . The primitive ray generators satisfy the relation

$$-\mathbf{e}_{E \setminus \text{cl}(I)} + \sum_{i \in F \setminus \text{cl}(I)} \mathbf{e}_i = -\mathbf{e}_{E \setminus F},$$

which is zero modulo the span of $\sigma_{I \leq \mathcal{F}}$. As the \mathbf{e}_i 's are independent modulo the span of $\sigma_{I \leq \mathcal{F}}$, any relation between the primitive generators must be a multiple of the displayed one.

When the closure of I is F , the fact that $\sigma_{I \leq \mathcal{F}}$ has codimension 1 implies that there is a unique integer k with $\text{rk } F < k < \text{rk } M$ such that \mathcal{F} does not include a flat of rank k . Let F_\circ be the unique flat in \mathcal{F} of rank $k - 1$, and let F° be the unique flat in $\mathcal{F} \cup \{E\}$ of rank $k + 1$. The primitive ray generators in question are $-\mathbf{e}_{E \setminus G}$ for the flats G in \mathcal{G} , where \mathcal{G} is the set of flats of M covering F_\circ and covered by F° . By the flat partition property of matroids [Oxl11, Section 1.4], the primitive ray generators satisfy the relation

$$\sum_{G \in \mathcal{G}} -\mathbf{e}_{E \setminus G} = -(\ell - 1)\mathbf{e}_{E \setminus F_\circ} - \mathbf{e}_{E \setminus F^\circ},$$

which is zero modulo the span of $\sigma_{I \leq \mathcal{F}}$. Since any proper subset of the primitive generators $-\mathbf{e}_{E \setminus G}$ for G in \mathcal{G} is independent modulo the span of $\sigma_{I \leq \mathcal{F}}$, any relation between the primitive generators must be a multiple of the displayed one.

The local analysis above shows that any constant d -dimensional weight on Π_M is balanced. Since Π_M is connected in codimension 1 by Proposition 2.3, it also shows that any d -dimensional balanced weight on Π_M must be constant. \square

2.4. Chow rings. Any unimodular fan Σ in \mathbb{R}^E defines a graded commutative algebra $\text{CH}(\Sigma)$, which is the Chow ring of the associated smooth toric variety X_Σ over \mathbb{C} with rational coefficients. Equivalently, $\text{CH}(\Sigma)$ is the ring of continuous piecewise polynomial functions on Σ with rational coefficients modulo the ideal generated by globally linear functions [Bri96, Section 3.1]. We write $\text{CH}^k(\Sigma)$ for the Chow group of codimension k cycles in X_Σ , so that

$$\text{CH}(\Sigma) = \bigoplus_k \text{CH}^k(\Sigma).$$

The group of k -dimensional balanced weights on Σ is related to $\text{CH}^k(\Sigma)$ by the isomorphism

$$\text{MW}_k(\Sigma) \longrightarrow \text{Hom}_{\mathbb{Q}}(\text{CH}^k(\Sigma), \mathbb{Q}), \quad \omega \longmapsto (x_\sigma \longmapsto \omega(\sigma)),$$

where x_σ is the class of the torus orbit closure in X_Σ corresponding to a k -dimensional cone σ in Σ . See [AHK18, Section 5] for a detailed discussion. For general facts on toric varieties and Chow rings, and for any undefined terms, we refer to [CLS11] and [Ful98].

In Proposition 2.10 below, we show that the Chow ring of M coincides with $\text{CH}(\underline{\Pi}_M)$ and that the augmented Chow ring of M coincides with $\text{CH}(\Pi_M)$.

Lemma 2.9. The following identities hold in the augmented Chow ring $\text{CH}(M)$.

- (1) For any element i of E , we have $y_i^2 = 0$.
- (2) For any two bases I_1 and I_2 of a flat F of M , we have $\prod_{i \in I_1} y_i = \prod_{i \in I_2} y_i$.
- (3) For any dependent set J of M , we have $\prod_{j \in J} y_j = 0$.

Proof. The first identity is a straightforward consequence of the relations in I_M and J_M :

$$y_i^2 = y_i \left(\sum_{i \notin F} x_F \right) = 0.$$

For the second identity, we may assume that $I_1 \setminus I_2 = \{i_1\}$ and $I_2 \setminus I_1 = \{i_2\}$, by the basis exchange property of matroids. Since a flat of M contains I_1 if and only if it contains I_2 , we have

$$\left(\sum_{i_1 \in G} x_G \right) \prod_{i \in I_1 \cap I_2} y_i = \left(\sum_{I_1 \subseteq G} x_G \right) \prod_{i \in I_1 \cap I_2} y_i = \left(\sum_{I_2 \subseteq G} x_G \right) \prod_{i \in I_1 \cap I_2} y_i = \left(\sum_{i_2 \in G} x_G \right) \prod_{i \in I_1 \cap I_2} y_i.$$

This immediately implies that we also have

$$\left(\sum_{i_1 \notin G} x_G \right) \prod_{i \in I_1 \cap I_2} y_i = \left(\sum_{i_2 \notin G} x_G \right) \prod_{i \in I_1 \cap I_2} y_i,$$

which tells us that

$$\prod_{i \in I_1} y_i = y_{i_1} \prod_{i \in I_1 \cap I_2} y_i = \left(\sum_{i_1 \notin G} x_G \right) \prod_{i \in I_1 \cap I_2} y_i = \left(\sum_{i_2 \notin G} x_G \right) \prod_{i \in I_1 \cap I_2} y_i = y_{i_2} \prod_{i \in I_1 \cap I_2} y_i = \prod_{i \in I_2} y_i.$$

For the third identity, we may suppose that J is a circuit, that is, a minimal dependent set. Since M is a loopless matroid, we may choose distinct elements j_1 and j_2 from J . Note that the independent sets $J \setminus j_1$ and $J \setminus j_2$ have the same closure because J is a circuit. Therefore, by the second identity, we have

$$\prod_{j \in J \setminus j_1} y_j = \prod_{j \in J \setminus j_2} y_j.$$

Combining the above with the first identity, we get

$$\prod_{j \in J} y_j = y_{j_1} \prod_{j \in J \setminus j_1} y_j = y_{j_1} \prod_{j \in J \setminus j_2} y_j = y_{j_1}^2 \prod_{j \in J \setminus \{j_1, j_2\}} y_j = 0. \quad \square$$

By the second identity in Lemma 2.9, we may define

$$y_F := \prod_{i \in I} y_i \text{ in } \text{CH}(M)$$

for any flat F of M and any basis I of F . The element y_E will play the role of the fundamental class for the augmented Chow ring of M .

Proposition 2.10. We have isomorphisms

$$\underline{\text{CH}}(M) \cong \text{CH}(\underline{\Pi}_M) \quad \text{and} \quad \text{CH}(M) \cong \text{CH}(\Pi_M).$$

Proof. The first isomorphism is proved in [FY04, Theorem 3]; see also [AHK18, Section 5.3].

Let K_M be the ideal of S_M generated by the monomials $\prod_{j \in J} y_j$ for every dependent set J of M . The ring of continuous piecewise polynomial functions on Π_M is isomorphic to the Stanley–Reisner ring of Δ_M , which is equal to

$$S_M / (J_M + K_M).$$

The ring $\text{CH}(\Pi_M)$ is obtained from this ring by killing the linear forms that generate the ideal I_M . In other words, we have a surjective homomorphism

$$\text{CH}(M) := S_M / (I_M + J_M) \longrightarrow S_M / (I_M + J_M + K_M) \cong \text{CH}(\Pi_M).$$

The fact that this is an isomorphism follows from the third part of Lemma 2.9. □

Remark 2.11. By Proposition 2.10, the graded dimension of the Chow ring of the rank d Boolean matroid $\underline{\text{CH}}(\text{B})$ is given by the h -vector of the permutohedron in \mathbb{R}^E . In other words, we have

$$\dim \underline{\text{CH}}^k(\text{B}) = \text{the Eulerian number } \left\langle \begin{matrix} d \\ k \end{matrix} \right\rangle.$$

See [Pet15, Section 9.1] for more on permutohedra and Eulerian numbers.

If E is nonempty, we have the balanced weight

$$1 \in \text{MW}_{d-1}(\underline{\Pi}_M) \cong \text{Hom}_{\mathbb{Q}}(\underline{\text{CH}}^{d-1}(M), \mathbb{Q}),$$

which can be used to define a degree map on the Chow ring of M . Similarly, for any E ,

$$1 \in \text{MW}_d(\Pi_M) \cong \text{Hom}_{\mathbb{Q}}(\text{CH}^d(M), \mathbb{Q})$$

can be used to define a degree map on the augmented Chow ring of M .

Definition 2.12. Consider the following *degree maps* for the Chow ring and the augmented Chow ring of M .

(1) If E is nonempty, the degree map for $\underline{\text{CH}}(M)$ is the linear map

$$\underline{\text{deg}}_M : \underline{\text{CH}}^{d-1}(M) \longrightarrow \mathbb{Q}, \quad x_{\mathcal{F}} \longmapsto 1,$$

where $x_{\mathcal{F}}$ is any monomial corresponding to a maximal cone $\underline{\sigma}_{\mathcal{F}}$ of $\underline{\Pi}_M$.

(2) For any E , the degree map for $\text{CH}(M)$ is the linear map

$$\text{deg}_M : \text{CH}^d(M) \longrightarrow \mathbb{Q}, \quad x_{I \leq \mathcal{F}} \longmapsto 1,$$

where $x_{I \leq \mathcal{F}}$ is any monomial corresponding to a maximal cone $\sigma_{I \leq \mathcal{F}}$ of Π_M .

By Proposition 2.8, the degree maps are well-defined and are isomorphisms. It follows that, for any two maximal cones $\underline{\sigma}_{\mathcal{F}_1}$ and $\underline{\sigma}_{\mathcal{F}_2}$ of the Bergman fan of M ,

$$x_{\mathcal{F}_1} = x_{\mathcal{F}_2} \text{ in } \underline{\text{CH}}^{d-1}(M).$$

Similarly, for any two maximal cones $\sigma_{I_1 \leq \mathcal{F}_1}$ and $\sigma_{I_2 \leq \mathcal{F}_2}$ of the augmented Bergman fan of M ,

$$y_{F_1} x_{\mathcal{F}_1} = y_{F_2} x_{\mathcal{F}_2} \text{ in } \text{CH}^d(M),$$

where F_1 is the closure of I_1 in M and F_2 is the closure of I_2 in M . Proposition 2.10 shows that

$$\underline{\text{CH}}^k(M) = 0 \text{ for } k \geq d \text{ and } \text{CH}^k(M) = 0 \text{ for } k > d.$$

Remark 2.13. Let \mathbb{F} be a field, and let V be a d -dimensional linear subspace of \mathbb{F}^E . We suppose that the subspace V is not contained in $\mathbb{F}^S \subseteq \mathbb{F}^E$ for any proper subset S of E . Let B be the Boolean matroid on E , and let M be the loopless matroid on E defined by

S is an independent set of $M \iff$ the restriction to V of the projection $\mathbb{F}^E \rightarrow \mathbb{F}^S$ is surjective.

Let $\mathbb{P}(\mathbb{F}^E)$ be the projective space of lines in \mathbb{F}^E , and let \mathbb{T}_E be its open torus. For any proper flat F of M , we write H_F for the projective subspace

$$H_F := \{p \in \mathbb{P}(V) \mid p_i = 0 \text{ for all } i \in F\}.$$

The *wonderful variety* \underline{X}_V is obtained from $\mathbb{P}(V)$ by first blowing up H_F for every corank 1 flat F , then blowing up the strict transforms of H_F for every corank 2 flat F , and so on. Equivalently,

$$\begin{aligned} \underline{X}_V &= \text{the closure of } \mathbb{P}(V) \cap \mathbb{T}_E \text{ in the toric variety } \underline{X}_M \text{ defined by } \Pi_M \\ &= \text{the closure of } \mathbb{P}(V) \cap \mathbb{T}_E \text{ in the toric variety } \underline{X}_B \text{ defined by } \Pi_B. \end{aligned}$$

When E is nonempty, the inclusion $\underline{X}_V \subseteq \underline{X}_M$ induces an isomorphism between their Chow rings,⁸ and hence the Chow ring of \underline{X}_V is isomorphic to $\underline{\text{CH}}(M)$ [FY04, Corollary 2].

Let $\mathbb{P}(\mathbb{F}^E \oplus \mathbb{F})$ be the projective completion of \mathbb{F}^E , and let \mathbb{T}_E be its open torus. The projective completion $\mathbb{P}(V \oplus \mathbb{F})$ contains a copy of $\mathbb{P}(V)$ as the hyperplane at infinity, and it therefore contains a copy of H_F for every nonempty proper flat F . The *augmented wonderful variety* X_V is obtained from $\mathbb{P}(V \oplus \mathbb{F}^1)$ by first blowing up H_F for every corank 1 flat F , then blowing up the strict transforms of H_F for every corank 2 flat F , and so on. Equivalently,

$$\begin{aligned} X_V &= \text{the closure of } \mathbb{P}(V \oplus \mathbb{F}) \cap \mathbb{T}_E \text{ in the toric variety } X_M \text{ defined by } \Pi_M \\ &= \text{the closure of } \mathbb{P}(V \oplus \mathbb{F}) \cap \mathbb{T}_E \text{ in the toric variety } X_B \text{ defined by } \Pi_B. \end{aligned}$$

The inclusion $X_V \subseteq X_M$ induces an isomorphism between their Chow rings, and hence the Chow ring of X_V is isomorphic to $\text{CH}(M)$.⁹

2.5. The graded Möbius algebra. For any nonnegative integer k , we define a vector space

$$H^k(M) := \bigoplus_{F \in \mathcal{L}^k(M)} \mathbb{Q}y_F,$$

where the direct sum is over the set $\mathcal{L}^k(M)$ of rank k flats of M .

Definition 2.14. The *graded Möbius algebra* of M is the graded vector space

$$H(M) := \bigoplus_{k \geq 0} H^k(M).$$

The multiplication in $H(M)$ is defined by the rule

$$y_{F_1}y_{F_2} = \begin{cases} y_{F_1 \vee F_2} & \text{if } \text{rk}_M(F_1) + \text{rk}_M(F_2) = \text{rk}_M(F_1 \vee F_2), \\ 0 & \text{if } \text{rk}_M(F_1) + \text{rk}_M(F_2) > \text{rk}_M(F_1 \vee F_2), \end{cases}$$

where \vee stands for the join operation in the lattice of flats $\mathcal{L}(M)$ of M .

⁸In general, the inclusion $\underline{X}_V \subseteq \underline{X}_M$ does not induce an isomorphism between their singular cohomology rings.

⁹This can be proved using the interpretation of $\text{CH}(M)$ in the last sentence of Remark 4.1.

Our double use of the symbol y_F is justified by the following proposition.

Proposition 2.15. The graded linear map

$$H(M) \longrightarrow \text{CH}(M), \quad y_F \longmapsto y_F$$

is an injective homomorphism of graded algebras.

Proof. We first show that the linear map is injective. It is enough to check that the subset

$$\{y_F\}_{F \in \mathcal{L}^k(M)} \subseteq \text{CH}^k(M)$$

is linearly independent for every nonnegative integer $k < d$. Suppose that

$$\sum_{F \in \mathcal{L}^k(M)} c_F y_F = 0 \text{ for some } c_F \in \mathbb{Q}.$$

For any given rank k flat G , we choose a saturated flag of proper flats \mathcal{G} whose smallest member is G and observe that

$$c_G y_G x_{\mathcal{G}} = \left(\sum_{F \in \mathcal{L}^k(M)} c_F y_F \right) x_{\mathcal{G}} = 0.$$

Since the degree of $y_G x_{\mathcal{G}}$ is 1, this implies that c_G must be zero.

We next check that the linear map is an algebra homomorphism using Lemma 2.9. Let I_1 be a basis of a flat F_1 , and let I_2 be a basis of a flat F_2 . If the rank of $F_1 \vee F_2$ is the sum of the ranks of F_1 and F_2 , then I_1 and I_2 are disjoint and their union is a basis of $F_1 \vee F_2$. Therefore, in the augmented Chow ring of M ,

$$y_{F_1} y_{F_2} = \prod_{i \in I_1} y_i \prod_{i \in I_2} y_i = \prod_{i \in I_1 \cup I_2} y_i = y_{F_1 \vee F_2}.$$

If the rank of $F_1 \vee F_2$ is less than the sum of the ranks of F_1 and F_2 , then either I_1 and I_2 intersect or the union of I_1 and I_2 is dependent in M . Therefore, in the augmented Chow ring of M ,

$$y_{F_1} y_{F_2} = \prod_{i \in I_1} y_i \prod_{i \in I_2} y_i = 0. \quad \square$$

Remark 2.16. Consider the torus \mathbb{T}_E , the toric variety X_B , and the augmented wonderful variety X_V in Remark 2.13. The identity of \mathbb{T}_E uniquely extends to a toric map

$$p_B : X_B \longrightarrow (\mathbb{P}^1)^E.$$

Let p_V be the restriction of p_B to the augmented wonderful variety X_V . If we identify the Chow ring of X_V with $\text{CH}(M)$ as in Remark 2.13, the image of the pullback p_V^* is the graded Möbius algebra $H(M) \subseteq \text{CH}(M)$.

2.6. Pullback and pushforward maps. Let Σ be a unimodular fan, and let σ be a k -dimensional cone in Σ . The torus orbit closure in the smooth toric variety X_Σ corresponding to σ can be identified with the toric variety of the fan $\text{star}_\sigma \Sigma$. Its class in the Chow ring of X_Σ is the monomial x_σ , which is the product of the divisor classes x_ρ corresponding to the rays ρ in σ . The inclusion ι of the torus orbit closure in X_Σ defines the *pullback* ι^* and the *pushforward* ι_* between the Chow rings, whose composition is multiplication by the monomial x_σ :

$$\begin{array}{ccc} \text{CH}(\Sigma) & \xrightarrow{x_\sigma} & \text{CH}(\Sigma) \\ & \searrow \iota^* & \nearrow \iota_* \\ & \text{CH}(\text{star}_\sigma \Sigma) & \end{array}$$

The pullback ι^* is a surjective graded algebra homomorphism, while the pushforward ι_* is a degree k homomorphism of $\text{CH}(\Sigma)$ -modules.

We give an explicit description of the pullback ι^* and the pushforward ι_* when Σ is the augmented Bergman fan Π_M and σ is the ray ρ_F of a proper flat F of M . Recall from Proposition 2.7 that the star of ρ_F admits the decomposition

$$\text{star}_{\rho_F} \Pi_M \cong \underline{\Pi}_{M_F} \times \Pi_{M^F}.$$

Thus we may identify the Chow ring of the star of ρ_F with $\underline{\text{CH}}(M_F) \otimes \text{CH}(M^F)$. We denote the pullback to the tensor product by φ_M^F and the pushforward from the tensor product by ψ_M^F :

$$\begin{array}{ccc} \text{CH}(M) & \xrightarrow{x_F} & \text{CH}(M) \\ & \searrow \varphi_M^F & \nearrow \psi_M^F \\ & \underline{\text{CH}}(M_F) \otimes \text{CH}(M^F) & \end{array}$$

To describe the pullback and the pushforward, we introduce Chow classes α_M , $\underline{\alpha}_M$, and $\underline{\beta}_M$. They are defined as the sums

$$\alpha_M := \sum_G x_G \in \text{CH}^1(M),$$

where the sum is over all proper flats G of M ;

$$\underline{\alpha}_M := \sum_{i \in G} x_G \in \underline{\text{CH}}^1(M),$$

where the sum is over all nonempty proper flats G of M containing a given element i in E ; and

$$\underline{\beta}_M := \sum_{i \notin G} x_G \in \underline{\text{CH}}^1(M),$$

where the sum is over all nonempty proper flats G of M not containing a given element i in E . The linear relations defining $\underline{\text{CH}}(M)$ show that $\underline{\alpha}_M$ and $\underline{\beta}_M$ do not depend on the choice of i .

The following two propositions are straightforward.

Proposition 2.17. The pullback φ_M^F is the unique graded algebra homomorphism

$$\mathrm{CH}(M) \longrightarrow \underline{\mathrm{CH}}(M_F) \otimes \mathrm{CH}(M^F)$$

that satisfies the following properties:

- If G is a proper flat of M incomparable to F , then $\varphi_M^F(x_G) = 0$.
- If G is a proper flat of M properly contained in F , then $\varphi_M^F(x_G) = 1 \otimes x_G$.
- If G is a proper flat of M properly containing F , then $\varphi_M^F(x_G) = x_{G \setminus F} \otimes 1$.
- If i is an element of F , then $\varphi_M^F(y_i) = 1 \otimes y_i$.
- If i is an element of $E \setminus F$, then $\varphi_M^F(y_i) = 0$.

The above five properties imply the following additional properties of φ_M^F :

- The equality $\varphi_M^F(x_F) = -1 \otimes \alpha_{M^F} - \underline{\beta}_{M^F} \otimes 1$ holds.
- The equality $\varphi_M^F(\alpha_M) = \underline{\alpha}_{M^F} \otimes 1$ holds.

Proposition 2.18. The pushforward ψ_M^F is the unique $\mathrm{CH}(M)$ -module homomorphism¹⁰

$$\psi_M^F : \underline{\mathrm{CH}}(M_F) \otimes \mathrm{CH}(M^F) \longrightarrow \mathrm{CH}(M)$$

that satisfies, for any collection \mathcal{S}' of proper flats of M strictly containing F and any collection \mathcal{S}'' of proper flats of M strictly contained in F ,

$$\psi_M^F \left(\prod_{F' \in \mathcal{S}'} x_{F' \setminus F} \otimes \prod_{F'' \in \mathcal{S}''} x_{F''} \right) = x_F \prod_{F' \in \mathcal{S}'} x_{F'} \prod_{F'' \in \mathcal{S}''} x_{F''}.$$

The composition $\psi_M^F \circ \varphi_M^F$ is multiplication by the element x_F , and the composition $\varphi_M^F \circ \psi_M^F$ is multiplication by the element $\varphi_M^F(x_F)$.

Proposition 2.18 shows that the pushforward ψ_M^F commutes with the degree maps:

$$\underline{\mathrm{deg}}_{M_F} \otimes \mathrm{deg}_{M^F} = \mathrm{deg}_M \circ \psi_M^F.$$

Proposition 2.19. If $\underline{\mathrm{CH}}(M_F)$ and $\mathrm{CH}(M^F)$ satisfy the Poincaré duality part of Theorem 1.3, then ψ_M^F is injective.

In other words, assuming Poincaré duality for the Chow rings, the graded $\mathrm{CH}(M)$ -module $\underline{\mathrm{CH}}(M_F) \otimes \mathrm{CH}(M^F)[-1]$ is isomorphic to the principal ideal of x_F in $\mathrm{CH}(M)$.¹¹ In particular,

$$\underline{\mathrm{CH}}(M)[-1] \cong \mathrm{ideal}(x_{\emptyset}) \subseteq \mathrm{CH}(M).$$

¹⁰We make ψ_M^F into a $\mathrm{CH}(M)$ -module homomorphism via the pullback φ_M^F .

¹¹For a graded vector space V , we write $V[m]$ for the graded vector space whose degree k piece is equal to V^{k+m} .

Proof. We will use the symbol \deg_F to denote the degree function $\underline{\deg}_{M_F} \otimes \deg_{M^F}$. For contradiction, suppose that $\psi_M^F(\eta) = 0$ for $\eta \neq 0$. By the two Poincaré duality statements in Theorem 1.3, there is an element ν such that $\deg_F(\nu\eta) = 1$. By surjectivity of the pullback φ_M^F , there is an element μ such that $\nu = \varphi_M^F(\mu)$. Since ψ_M^F is a $\text{CH}(M)$ -module homomorphism that commutes with the degree maps, we have

$$1 = \deg_F(\nu\eta) = \deg_M(\psi_M^F(\nu\eta)) = \deg_M(\psi_M^F(\varphi_M^F(\mu)\eta)) = \deg_M(\mu\psi_M^F(\eta)) = \deg_M(0) = 0,$$

which is a contradiction. \square

We next give an explicit description of the pullback ι^* and the pushforward ι_* when Σ is the Bergman fan $\underline{\Pi}_M$ and σ is the ray ρ_F of a nonempty proper flat F of M . Recall from Proposition 2.7 that the star of ρ_F admits the decomposition

$$\text{star}_{\rho_F} \underline{\Pi}_M \cong \underline{\Pi}_{M_F} \times \underline{\Pi}_{M^F}.$$

Thus we may identify the Chow ring of the star of ρ_F with $\underline{\text{CH}}(M_F) \otimes \underline{\text{CH}}(M^F)$. We denote the pullback to the tensor product by φ_M^F and the pushforward from the tensor product by ψ_M^F :

$$\begin{array}{ccc} \underline{\text{CH}}(M) & \xrightarrow{x_F} & \underline{\text{CH}}(M) \\ & \searrow \varphi_M^F & \nearrow \psi_M^F \\ & \underline{\text{CH}}(M_F) \otimes \underline{\text{CH}}(M^F) & \end{array}$$

The following analogues of Propositions 2.17 and 2.18 are straightforward.

Proposition 2.20. The pullback φ_M^F is the unique graded algebra homomorphism

$$\underline{\text{CH}}(M) \longrightarrow \underline{\text{CH}}(M_F) \otimes \underline{\text{CH}}(M^F)$$

that satisfies the following properties:

- If G is a nonempty proper flat of M incomparable to F , then $\varphi_M^F(x_G) = 0$.
- If G is a nonempty proper flat of M properly contained in F , then $\varphi_M^F(x_G) = 1 \otimes x_G$.
- If G is a nonempty proper flat of M properly containing F , then $\varphi_M^F(x_G) = x_{G \setminus F} \otimes 1$.

The above three properties imply the following additional properties of φ_M^F :

- The equality $\varphi_M^F(x_F) = -1 \otimes \underline{\alpha}_{M^F} - \underline{\beta}_{M_F} \otimes 1$ holds.
- The equality $\varphi_M^F(\underline{\alpha}_M) = \underline{\alpha}_{M_F} \otimes 1$ holds.
- The equality $\varphi_M^F(\underline{\beta}_M) = 1 \otimes \underline{\beta}_{M^F}$ holds.

Proposition 2.21. The pushforward $\underline{\psi}_M^F$ is the unique $\underline{\text{CH}}(M)$ -module homomorphism

$$\underline{\text{CH}}(M_F) \otimes \underline{\text{CH}}(M^F) \longrightarrow \underline{\text{CH}}(M)$$

that satisfies, for any collection \mathcal{S}' of proper flats of M strictly containing F and any collection \mathcal{S}'' of nonempty proper flats of M strictly contained in F ,

$$\underline{\psi}_M^F \left(\prod_{F' \in \mathcal{S}'} x_{F' \setminus F} \otimes \prod_{F'' \in \mathcal{S}''} x_{F''} \right) = x_F \prod_{F' \in \mathcal{S}'} x_{F'} \prod_{F'' \in \mathcal{S}''} x_{F''}.$$

The composition $\underline{\psi}_M^F \circ \underline{\varphi}_M^F$ is multiplication by the element x_F , and the composition $\underline{\varphi}_M^F \circ \underline{\psi}_M^F$ is multiplication by the element $\underline{\varphi}_M^F(x_F)$.

Proposition 2.21 shows that the pushforward $\underline{\psi}_M^F$ commutes with the degree maps:

$$\underline{\text{deg}}_{M_F} \otimes \underline{\text{deg}}_{M^F} = \underline{\text{deg}}_M \circ \underline{\psi}_M^F.$$

Proposition 2.22. If $\underline{\text{CH}}(M_F)$ and $\underline{\text{CH}}(M^F)$ satisfy the Poincaré duality part of Theorem 1.3, then $\underline{\psi}_M^F$ is injective.

In other words, assuming Poincaré duality for the Chow rings, the graded $\underline{\text{CH}}(M)$ -module $\underline{\text{CH}}(M_F) \otimes \underline{\text{CH}}(M^F)[-1]$ is isomorphic to the principal ideal of x_F in $\underline{\text{CH}}(M)$.

Proof. The proof is essentially identical to that of Proposition 2.19. □

Last, we give an explicit description of the pullback ι^* and the pushforward ι_* when Σ is the augmented Bergman fan Π_M and σ is the cone σ_I of a nonempty independent set I of M . By Proposition 2.7, we have

$$\text{star}_{\sigma_I} \Pi_M \cong \Pi_{M_F},$$

where F is the closure of I in M . Thus we may identify the Chow ring of the star of σ_I with $\text{CH}(M_F)$. We denote the corresponding pullback by φ_F^M and the pushforward by ψ_F^M :

$$\begin{array}{ccc} \text{CH}(M) & \xrightarrow{y_F} & \text{CH}(M) \\ & \searrow \varphi_F^M & \nearrow \psi_F^M \\ & \text{CH}(M_F) & \end{array}$$

Note that the pullback and the pushforward only depend on F and not on I .

The following analogues of Propositions 2.17 and 2.18 are straightforward.

Proposition 2.23. The pullback φ_F^M is the unique graded algebra homomorphism

$$\text{CH}(M) \longrightarrow \text{CH}(M_F)$$

that satisfies the following properties:

- If G is a proper flat of M that contains F , then $\varphi_F^M(x_G) = x_{G \setminus F}$.
- If G is a proper flat of M that does not contain F , then $\varphi_F^M(x_G) = 0$.

The above two properties imply the following additional properties of φ_F^M :

- If i is an element of F , then $\varphi_F^M(y_i) = 0$.
- If i is an element of $E \setminus F$, then $\varphi_F^M(y_i) = y_i$.
- The equality $\varphi_F^M(\alpha_M) = \alpha_{M_F}$ holds.

Proposition 2.24. The pushforward ψ_F^M is the unique $\text{CH}(M)$ -module homomorphism

$$\text{CH}(M_F) \longrightarrow \text{CH}(M)$$

that satisfies, for any collection \mathcal{S}' of proper flats of M containing F ,

$$\psi_F^M \left(\prod_{F' \in \mathcal{S}'} x_{F' \setminus F} \right) = y_F \prod_{F' \in \mathcal{S}'} x_{F'}.$$

The composition $\psi_F^M \circ \varphi_F^M$ is multiplication by the element y_F , and the composition $\varphi_F^M \circ \psi_F^M$ is multiplication by the element $\varphi_F^M(y_F)$.

Proposition 2.24 shows that the pushforward ψ_F^M commutes with the degree maps:

$$\text{deg}_{M_F} = \text{deg}_M \circ \psi_F^M.$$

Proposition 2.25. If $\text{CH}(M_F)$ satisfies the Poincaré duality part of Theorem 1.3, then ψ_F^M is injective.

In other words, assuming Poincaré duality for the Chow rings, the graded $\text{CH}(M)$ -module $\text{CH}(M_F)[- \text{rk}_M(F)]$ is isomorphic to the principal ideal of y_F in $\text{CH}(M)$.

Proof. The proof is essentially identical to that of Proposition 2.19. □

The basic properties of the pullback and the pushforward maps can be used to describe the fundamental classes of $\underline{\text{CH}}(M)$ and $\text{CH}(M)$ in terms of $\underline{\alpha}_M$ and α_M .

Proposition 2.26. The degree of $\underline{\alpha}_M^{d-1}$ is 1, and the degree of α_M^d is 1.

Proof. We prove the first statement by induction on $d \geq 1$. Note that, for any nonempty proper flat F of rank k , we have

$$x_F \underline{\alpha}_M^{d-k} = \underline{\psi}_M^F(\underline{\varphi}_M^F(\underline{\alpha}_M^{d-k})) = \underline{\psi}_M^F(\underline{\alpha}_{M_F}^{d-k} \otimes 1) = 0,$$

since $\underline{\text{CH}}^{d-k}(M_F) = 0$. Therefore, for any proper flat a of rank 1 and any element i in a , we have

$$\underline{\alpha}_M^{d-1} = \left(\sum_{i \in F} x_F \right) \underline{\alpha}_M^{d-2} = x_a \underline{\alpha}_M^{d-2}.$$

Now, using the induction hypothesis applied to the matroid M_a of rank $d-1$, we get

$$\underline{\alpha}_M^{d-1} = x_a \underline{\alpha}_M^{d-2} = \underline{\psi}_M^a(\underline{\varphi}_M^a(\underline{\alpha}_M^{d-2})) = \underline{\psi}_M^a(\underline{\alpha}_{M_a}^{d-2} \otimes 1) = x_{\mathcal{F}},$$

where \mathcal{F} is any maximal flag of nonempty proper flats of M that starts from a .

For the second statement, note that, for any proper flat F of rank k ,

$$x_F \alpha_M^{d-k} = \psi_M^F(\varphi_M^F(\alpha_M^{d-k})) = \psi_M^F(\underline{\alpha}_{M_F}^{d-k} \otimes 1) = 0.$$

Using the first statement, we get the conclusion from the identity

$$\alpha_M^d = \left(\sum_F x_F \right) \alpha_M^{d-1} = x_{\emptyset} \alpha_M^{d-1} = \psi_M^{\emptyset}(\varphi_M^{\emptyset}(\alpha_M^{d-1})) = \psi_M^{\emptyset}(\underline{\alpha}_M^{d-1}). \quad \square$$

More generally, the degree of $\underline{\alpha}_M^{d-k} \underline{\beta}_M^k$ is the k -th coefficient of the reduced characteristic polynomial of M [AHK18, Proposition 9.5].

Remark 2.27. In the setting of Remark 2.13, the element α_M , viewed as an element of the Chow ring of the augmented wonderful variety X_V , is the class of the pullback of the hyperplane $\mathbb{P}(V) \subseteq \mathbb{P}(V \oplus \mathbb{F})$.

3. PROOFS OF THE SEMI-SMALL DECOMPOSITIONS AND THE POINCARÉ DUALITY THEOREMS

In this section, we prove Theorems 1.1 and 1.2 together with the two Poincaré duality statements in Theorem 1.3. For an element i of E , we write π_i and $\underline{\pi}_i$ for the coordinate projections

$$\pi_i : \mathbb{R}^E \longrightarrow \mathbb{R}^{E \setminus i} \quad \text{and} \quad \underline{\pi}_i : \mathbb{R}^E / \langle \mathbf{e}_E \rangle \longrightarrow \mathbb{R}^{E \setminus i} / \langle \mathbf{e}_{E \setminus i} \rangle.$$

Note that $\pi_i(\rho_i) = 0$ and $\underline{\pi}_i(\underline{\rho}_{\{i\}}) = 0$. In addition, $\pi_i(\rho_S) = \rho_{S \setminus i}$ and $\underline{\pi}_i(\underline{\rho}_S) = \underline{\rho}_{S \setminus i}$ for $S \subseteq E$.

Proposition 3.1. Let M be a loopless matroid on E , and let i be an element of E .

- (1) The projection π_i maps any cone of Π_M onto a cone of $\Pi_{M \setminus i}$.
- (2) The projection $\underline{\pi}_i$ maps any cone of $\underline{\Pi}_M$ onto a cone of $\underline{\Pi}_{M \setminus i}$.

Recall that a linear map defines a morphism of fans $\Sigma_1 \rightarrow \Sigma_2$ if it maps any cone of Σ_1 into a cone of Σ_2 [CLS11, Chapter 3]. Thus the above proposition is stronger than the statement that π_i and $\underline{\pi}_i$ induce morphisms of fans.

Proof. The projection π_i maps $\sigma_{I \leq \mathcal{F}}$ onto $\sigma_{I \setminus i \leq \mathcal{F} \setminus i}$, where $\mathcal{F} \setminus i$ is the flag of flats of $M \setminus i$ obtained by removing i from the members of \mathcal{F} . Similarly, $\underline{\pi}_i$ maps $\underline{\sigma}_{\mathcal{F}}$ onto $\underline{\sigma}_{\mathcal{F} \setminus i}$. \square

By Proposition 3.1, the projection π_i defines a map from the toric variety X_M of Π_M to the toric variety $X_{M \setminus i}$ of $\Pi_{M \setminus i}$, and hence the pullback homomorphism $\text{CH}(M \setminus i) \rightarrow \text{CH}(M)$. Explicitly, the pullback is the graded algebra homomorphism

$$\theta_i = \theta_i^M : \text{CH}(M \setminus i) \longrightarrow \text{CH}(M), \quad x_F \longmapsto x_F + x_{F \cup i},$$

where a variable in the target is set to zero if its label is not a flat of M . Similarly, $\underline{\pi}_i$ defines a map from the toric variety \underline{X}_M of $\underline{\Pi}_M$ to the toric variety $\underline{X}_{M \setminus i}$ of $\underline{\Pi}_{M \setminus i}$, and hence an algebra homomorphism

$$\underline{\theta}_i = \underline{\theta}_i^M : \underline{\text{CH}}(M \setminus i) \longrightarrow \underline{\text{CH}}(M), \quad x_F \longmapsto x_F + x_{F \cup i},$$

where a variable in the target is set to zero if its label is not a flat of M .

Remark 3.2. We use the notations introduced in Remark 2.13. Let $V \setminus i$ be the image of V under the i -th projection $\mathbb{F}^E \rightarrow \mathbb{F}^{E \setminus i}$. We have the commutative diagrams of wonderful varieties and their Chow rings

$$\begin{array}{ccc} \underline{X}_B & \longleftarrow & \underline{X}_V \\ \underline{p}_i^B \downarrow & & \downarrow \underline{p}_i^V \\ \underline{X}_{B \setminus i} & \longleftarrow & \underline{X}_{V \setminus i} \end{array} \qquad \begin{array}{ccc} \underline{\text{CH}}(B) & \longrightarrow & \underline{\text{CH}}(M) \\ \uparrow & & \uparrow \\ \underline{\text{CH}}(B \setminus i) & \longrightarrow & \underline{\text{CH}}(M \setminus i). \end{array}$$

The map \underline{p}_i^V is birational if and only if i is not a coloop of M . By Proposition 3.1, the fibers of \underline{p}_i^B are at most one-dimensional, and hence the fibers of \underline{p}_i^V are at most one-dimensional. It follows that \underline{p}_i^V is semi-small in the sense of Goresky–MacPherson when i is not a coloop of M .

Similarly, we have the diagrams of augmented wonderful varieties and their Chow rings

$$\begin{array}{ccc} X_B & \longleftarrow & X_V \\ p_i^B \downarrow & & \downarrow p_i^V \\ X_{B \setminus i} & \longleftarrow & X_{V \setminus i} \end{array} \qquad \begin{array}{ccc} \text{CH}(B) & \longrightarrow & \text{CH}(M) \\ \uparrow & & \uparrow \\ \text{CH}(B \setminus i) & \longrightarrow & \text{CH}(M \setminus i). \end{array}$$

The map p_i^V is birational if and only if i is not a coloop of M . By Proposition 3.1, the fibers of p_i^B are at most one-dimensional, and hence p_i^V is semi-small when i is not a coloop of M .

Numerically, the semi-smallness of \underline{p}_i^V is reflected in the identity

$$\dim x_{F \cup i} \underline{\text{CH}}_{(i)}^{k-1} = \dim x_{F \cup i} \underline{\text{CH}}_{(i)}^{d-k-2}.$$

Similarly, the semi-smallness of p_i^V is reflected in the identity¹²

$$\dim x_{F \cup i} \text{CH}_{(i)}^{k-1} = \dim x_{F \cup i} \text{CH}_{(i)}^{d-k-1}.$$

¹²The displayed identities follow from Proposition 3.5 and the Poincaré duality parts of Theorem 1.3.

For a detailed discussion of semi-small maps in the context of Hodge theory and the decomposition theorem, see [dCM02] and [dCM09].

The element i is said to be a *coloop* of M if the ranks of M and $M \setminus i$ are not equal. We show that the pullbacks θ_i and $\underline{\theta}_i$ are compatible with the degree maps of M and $M \setminus i$.

Lemma 3.3. Suppose that $E \setminus i$ is nonempty.

(1) If i is not a coloop of M , then θ_i commutes with the degree maps:

$$\deg_{M \setminus i} = \deg_M \circ \theta_i.$$

(2) If i is not a coloop of M , then $\underline{\theta}_i$ commutes with the degree maps:

$$\underline{\deg}_{M \setminus i} = \underline{\deg}_M \circ \underline{\theta}_i.$$

(3) If i is a coloop of M , we have

$$\deg_{M \setminus i} = \deg_M \circ x_{E \setminus i} \circ \theta_i = \deg_M \circ \alpha_M \circ \theta_i,$$

where the middle maps are multiplications by the elements $x_{E \setminus i}$ and α_M .

(4) If i is a coloop of M , we have

$$\underline{\deg}_{M \setminus i} = \underline{\deg}_M \circ x_{E \setminus i} \circ \underline{\theta}_i = \underline{\deg}_M \circ \underline{\alpha}_M \circ \underline{\theta}_i,$$

where the middle maps are multiplications by the elements $x_{E \setminus i}$ and $\underline{\alpha}_M$.

Proof. If i is not a coloop of M , we may choose a basis B of $M \setminus i$ that is also a basis of M . We have

$$\mathrm{CH}^d(M \setminus i) = \mathrm{span}(y_B) \quad \text{and} \quad \mathrm{CH}^d(M) = \mathrm{span}(y_B).$$

Since $\theta_i(y_j) = y_j$ for all j , the first identity follows. Similarly, by Proposition 2.26,

$$\underline{\mathrm{CH}}^{d-1}(M \setminus i) = \mathrm{span}(\underline{\alpha}_{M \setminus i}^{d-1}) \quad \text{and} \quad \underline{\mathrm{CH}}^{d-1}(M) = \mathrm{span}(\underline{\alpha}_M^{d-1}).$$

Since $\underline{\theta}_i(\underline{\alpha}_{M \setminus i}) = \underline{\alpha}_M$ when i is not a coloop, the second identity follows.

Suppose now that i is a coloop of M . In this case, $M \setminus i = M^{E \setminus i}$, and hence

$$\varphi_M^{E \setminus i} \circ \theta_i = \text{identity of } \mathrm{CH}(M \setminus i) \quad \text{and} \quad \underline{\varphi}_M^{E \setminus i} \circ \underline{\theta}_i = \text{identity of } \underline{\mathrm{CH}}(M \setminus i).$$

Using the compatibility of the pushforward $\psi_M^{E \setminus i}$ with the degree maps, we have

$$\deg_{M \setminus i} = \deg_M \circ \psi_M^{E \setminus i} = \deg_M \circ \psi_M^{E \setminus i} \circ \varphi_M^{E \setminus i} \circ \theta_i = \deg_M \circ x_{E \setminus i} \circ \theta_i.$$

Since $\theta_i(\alpha_{M \setminus i}) = \alpha_M - x_{E \setminus i}$ when i is a coloop of M , the above implies

$$\deg_{M \setminus i} = \deg_M \circ x_{E \setminus i} \circ \theta_i = \deg_M \circ (\alpha_M - \theta_i(\alpha_{M \setminus i})) \circ \theta_i = \deg_M \circ \alpha_M \circ \theta_i,$$

The identities for $\underline{\deg}_{M \setminus i}$ can be obtained in a similar way. □

Proposition 3.4. If $\text{CH}(M \setminus i)$ satisfies the Poincaré duality part of Theorem 1.3, then θ_i is injective. Also, if $\underline{\text{CH}}(M \setminus i)$ satisfies the Poincaré duality part of Theorem 1.3, then $\underline{\theta}_i$ is injective.

Proof. The proof is essentially identical to that of Proposition 2.19. \square

For a flat F in \mathcal{S}_i , we write $\theta_i^{F \cup i}$ for the pullback map between the augmented Chow rings obtained from the deletion of i from the localization $M^{F \cup i}$:

$$\theta_i^{F \cup i} : \text{CH}(M^F) \rightarrow \text{CH}(M^{F \cup i}).$$

Similarly, for a flat F in $\underline{\mathcal{S}}_i$, we write $\underline{\theta}_i^{F \cup i}$ for the pullback map between the Chow rings obtained from the deletion of i from the localization $M^{F \cup i}$:

$$\underline{\theta}_i^{F \cup i} : \underline{\text{CH}}(M^F) \rightarrow \underline{\text{CH}}(M^{F \cup i}).$$

Note that i is a coloop of $M^{F \cup i}$ in these cases.

Proposition 3.5. The summands appearing in Theorems 1.1 and 1.2 can be described as follows.

- (1) If $F \in \mathcal{S}_i$, then $x_{F \cup i} \text{CH}(i) = \psi_M^{F \cup i} (\underline{\text{CH}}(M_{F \cup i}) \otimes \theta_i^{F \cup i} \text{CH}(M^F))$.
- (2) If $F \in \underline{\mathcal{S}}_i$, then $x_{F \cup i} \underline{\text{CH}}(i) = \underline{\psi}_M^{F \cup i} (\underline{\text{CH}}(M_{F \cup i}) \otimes \underline{\theta}_i^{F \cup i} \underline{\text{CH}}(M^F))$.
- (3) If i is a coloop of M , then $x_{E \setminus i} \text{CH}(i) = \psi_M^{E \setminus i} \text{CH}(M \setminus i)$ and $x_{E \setminus i} \underline{\text{CH}}(i) = \underline{\psi}_M^{E \setminus i} \underline{\text{CH}}(M \setminus i)$.

It follows, assuming Poincaré duality for the Chow rings,¹³ that

$$x_{F \cup i} \text{CH}(i) \cong \underline{\text{CH}}(M_{F \cup i}) \otimes \text{CH}(M^F)[-1] \quad \text{and} \quad x_{F \cup i} \underline{\text{CH}}(i) \cong \underline{\text{CH}}(M_{F \cup i}) \otimes \underline{\text{CH}}(M^F)[-1].$$

Therefore, again assuming Poincaré duality for the Chow rings, we have

$$\dim x_{F \cup i} \underline{\text{CH}}(i)^{k-1} = \dim x_{F \cup i} \underline{\text{CH}}(i)^{d-k-2} \quad \text{and} \quad \dim x_{F \cup i} \text{CH}(i)^{k-1} = \dim x_{F \cup i} \text{CH}(i)^{d-k-1}.$$

Proof. We prove the first statement. The proof of the second statement is essentially identical. The third statement is a straightforward consequence of the fact that $\varphi_M^{E \setminus i} \circ \theta_i$ and $\underline{\varphi}_M^{E \setminus i} \circ \underline{\theta}_i$ are the identity maps when i is a coloop.

Let F be a flat in \mathcal{S}_i . It is enough to show that

$$\varphi_M^{F \cup i} (\text{CH}(i)) = \underline{\text{CH}}(M_{F \cup i}) \otimes \theta_i^{F \cup i} \text{CH}(M^F),$$

¹³We need Poincaré duality for $\underline{\text{CH}}(M^F)$, $\text{CH}(M^F)$, $\underline{\text{CH}}(M^{F \cup i})$, $\text{CH}(M^{F \cup i})$, and $\underline{\text{CH}}(M_{F \cup i})$.

since the result will then follow by applying $\psi_M^{F \cup i}$. The projection π_i maps the ray $\rho_{F \cup i}$ to the ray ρ_F , and hence π_i defines morphisms of fans

$$\begin{array}{ccccc} \text{star}_{\rho_{F \cup i}} \Pi_M & \xleftarrow{\iota_{F \cup i}} & \underline{\Pi}_{M_{F \cup i}} \times \Pi_{M^{F \cup i}} & \xlongequal{\quad} & \underline{\Pi}_{(M/i)_F} \times \Pi_{M^{F \cup i}} \\ \pi'_i \downarrow & & \pi''_i \downarrow & & \pi'''_i \downarrow \\ \text{star}_{\rho_F} \Pi_{M \setminus i} & \xleftarrow{\iota_F} & \underline{\Pi}_{(M \setminus i)_F} \times \Pi_{(M \setminus i)^F} & \xlongequal{\quad} & \underline{\Pi}_{(M \setminus i)_F} \times \Pi_{M^F}, \end{array}$$

where $\iota_{F \cup i}$ and ι_F are the isomorphisms in Proposition 2.7. The main point is that the matroid $(M/i)_F$ is a quotient of $(M \setminus i)_F$. In other words, we have the inclusion of Bergman fans

$$\underline{\Pi}_{(M/i)_F} \subseteq \underline{\Pi}_{(M \setminus i)_F}.$$

Therefore, the morphism π'''_i admits the factorization

$$\underline{\Pi}_{(M/i)_F} \times \Pi_{M^{F \cup i}} \longrightarrow \underline{\Pi}_{(M/i)_F} \times \Pi_{M^F} \hookrightarrow \underline{\Pi}_{(M \setminus i)_F} \times \Pi_{M^F},$$

where the second map induces a surjective pullback map q between the Chow rings. By the equality $(M/i)_F = M_{F \cup i}$, we have the commutative diagram of pullback maps between the Chow rings

$$\begin{array}{ccc} \text{CH}(M \setminus i) & \xrightarrow{\theta_i} & \text{CH}(M) \\ \varphi_{M \setminus i}^F \downarrow & & \downarrow \varphi_M^{F \cup i} \\ \underline{\text{CH}}((M \setminus i)_F) \otimes \text{CH}((M \setminus i)^F) & \xrightarrow{q} \underline{\text{CH}}(M_{F \cup i}) \otimes \text{CH}(M^F) & \xrightarrow{1 \otimes \theta_i^{F \cup i}} \underline{\text{CH}}(M_{F \cup i}) \otimes \text{CH}(M^{F \cup i}). \end{array}$$

The conclusion follows from the surjectivity of the pullback maps $\varphi_{M \setminus i}^F$ and q . \square

Remark 3.6. Since i is a coloop in $M^{F \cup i}$ when $F \in \mathcal{S}_i$ or $F \in \underline{\mathcal{S}}_i$, Proposition 3.5 implies that

$$x_{F \cup i} \text{CH}_{(i)}^{d-1} = 0 \text{ for } F \in \mathcal{S}_i \text{ and } x_{F \cup i} \underline{\text{CH}}_{(i)}^{d-2} = 0 \text{ for } F \in \underline{\mathcal{S}}_i.$$

Proposition 3.7. The Poincaré pairing on the summands appearing in Theorems 1.1 and 1.2 can be described as follows.

(1) If $F \in \mathcal{S}_i$, then for any $\mu_1, \mu_2 \in \underline{\text{CH}}(M_{F \cup i}) \otimes \text{CH}(M^F)$ of complementary degrees,

$$\deg_M (\psi_M^{F \cup i} (1 \otimes \theta_i^{F \cup i}(\mu_1)) \cdot \psi_M^{F \cup i} (1 \otimes \theta_i^{F \cup i}(\mu_2))) = -\underline{\deg}_{M_{F \cup i}} \otimes \deg_{M^F}(\mu_1 \mu_2).$$

(2) If $F \in \underline{\mathcal{S}}_i$, then for any $\nu_1, \nu_2 \in \underline{\text{CH}}(M_{F \cup i}) \otimes \underline{\text{CH}}(M^F)$ of complementary degrees,

$$\underline{\deg}_M (\underline{\psi}_M^{F \cup i} (1 \otimes \underline{\theta}_i^{F \cup i}(\nu_1)) \cdot \underline{\psi}_M^{F \cup i} (1 \otimes \underline{\theta}_i^{F \cup i}(\nu_2))) = -\underline{\deg}_{M_{F \cup i}} \otimes \underline{\deg}_{M^F}(\nu_1 \nu_2).$$

It follows, assuming Poincaré duality for the Chow rings,¹⁴ that the restriction of the Poincaré pairing of $\text{CH}(M)$ to the subspace $x_{F \cup i} \text{CH}_{(i)}$ is non-degenerate, and the restriction of the Poincaré pairing of $\underline{\text{CH}}(M)$ to the subspace $x_{F \cup i} \underline{\text{CH}}_{(i)}$ is non-degenerate.

Proof. We prove the first identity. The second identity can be proved in the same way.

Since the pushforward $\psi_M^{F \cup i}$ is a $\text{CH}(M)$ -module homomorphism, the left-hand side is

$$\deg_M (\psi_M^{F \cup i} (\varphi_M^{F \cup i} \psi_M^{F \cup i} (1 \otimes \theta_i^{F \cup i}(\mu_1)) \cdot (1 \otimes \theta_i^{F \cup i}(\mu_2)))).$$

The pushforward commutes with the degree maps, so the above is equal to

$$\underline{\deg}_{M^{F \cup i}} \otimes \deg_{M^{F \cup i}} (\varphi_M^{F \cup i} \psi_M^{F \cup i} (1 \otimes \theta_i^{F \cup i}(\mu_1)) \cdot (1 \otimes \theta_i^{F \cup i}(\mu_2))).$$

Using that the composition $\varphi_M^{F \cup i} \psi_M^{F \cup i}$ is multiplication by $\varphi_M^{F \cup i}(x_{F \cup i})$, we get

$$-\underline{\deg}_{M^{F \cup i}} \otimes \deg_{M^{F \cup i}} ((1 \otimes \alpha_{M^{F \cup i}} + \beta_{M^{F \cup i}} \otimes 1) \cdot (1 \otimes \theta_i^{F \cup i}(\mu_1)) \cdot (1 \otimes \theta_i^{F \cup i}(\mu_2))).$$

Since i is a coloop of $M^{F \cup i}$, the expression simplifies to

$$-\underline{\deg}_{M^{F \cup i}} \otimes \deg_{M^{F \cup i}} ((1 \otimes \alpha_{M^{F \cup i}}) \cdot (1 \otimes \theta_i^{F \cup i}(\mu_1)) \cdot (1 \otimes \theta_i^{F \cup i}(\mu_2))).$$

Now the third part of Lemma 3.3 shows that the above quantity is the right-hand side of the formula in statement (1). \square

Lemma 3.8. If flats F_1, F_2 are in \mathcal{S}_i and F_1 is a proper subset of F_2 , then

$$x_{F_1 \cup i} x_{F_2 \cup i} \in x_{F_1 \cup i} \text{CH}_{(i)}.$$

Similarly, if F_1, F_2 are in $\underline{\mathcal{S}}_i$ and F_1 is a proper subset of F_2 , then

$$x_{F_1 \cup i} x_{F_2 \cup i} \in x_{F_1 \cup i} \underline{\text{CH}}_{(i)}.$$

Proof. Since $F_1 \cup i$ is not comparable to F_2 , we have

$$x_{F_1 \cup i} x_{F_2 \cup i} = x_{F_1 \cup i} (x_{F_2} + x_{F_2 \cup i}) = x_{F_1 \cup i} \theta_i(x_{F_2}).$$

The second part follows from the same argument. \square

Proof of Theorem 1.1, Theorem 1.2, and parts (1) and (4) of Theorem 1.3. All the summands in the proposed decompositions are cyclic, and therefore indecomposable in the category of graded modules.¹⁵ We prove the decompositions by induction on the cardinality of the ground set E . If E is empty, then Theorem 1.1, Theorem 1.2, and part (1) of Theorem 1.3 are vacuous, while part (4) of Theorem 1.3 is trivial. Furthermore, all of these results are trivial when E is a singleton. Thus,

¹⁴We need Poincaré duality for $\underline{\text{CH}}(M^F)$, $\text{CH}(M^F)$, $\underline{\text{CH}}(M^{F \cup i})$, $\text{CH}(M^{F \cup i})$, and $\underline{\text{CH}}(M^{F \cup i})$.

¹⁵By [CF82, Corollary 2] or [GG82, Theorem 3.2], the indecomposability of the summands in the category of graded modules implies the indecomposability of the summands in the category of modules.

we may assume that i is an element of E , that $E \setminus i$ is nonempty, and that all the results hold for loopless matroids whose ground set is a proper subset of E .

First we assume that i is not a coloop. Let us show that the terms in the right-hand side of the decomposition (D_1) are orthogonal. Multiplying $\text{CH}_{(i)}$ and $x_{F \cup i} \text{CH}_{(i)}$ lands in $x_{F \cup i} \text{CH}_{(i)}$, and this ideal vanishes in degree d by Remark 3.6, so they are orthogonal. On the other hand, the product of $x_{F_1 \cup i} \text{CH}_{(i)}$ and $x_{F_2 \cup i} \text{CH}_{(i)}$ vanishes if $F_1, F_2 \in \mathcal{S}_i$ are not comparable, while if $F_1 < F_2$ or $F_2 < F_1$, the product is contained in $x_{F_1 \cup i} \text{CH}_{(i)}$ or $x_{F_2 \cup i} \text{CH}_{(i)}$ respectively, by Lemma 3.8. So these terms are also orthogonal.

It follows from the induction hypothesis and Lemma 3.3 that the restriction of the Poincaré pairing of $\text{CH}(M)$ to $\text{CH}_{(i)}$ is non-degenerate. By Proposition 3.5, Proposition 3.7, and the induction hypothesis, the restriction of the Poincaré pairing of $\text{CH}(M)$ to any other summand $x_{F \cup i} \text{CH}_{(i)}$ is also non-degenerate. Therefore, we can conclude that the sum on the right-hand side of (D_1) is a direct sum with a non-degenerate Poincaré pairing.

To complete the proof of the decomposition (D_1) and the Poincaré duality theorem for $\text{CH}(M)$, we must show that the direct sum

$$\text{CH}_{(i)} \oplus \bigoplus_{F \in \mathcal{S}_i} x_{F \cup i} \text{CH}_{(i)}$$

is equal to all of $\text{CH}(M)$. This is obvious in degree 0. To see that it holds in degree 1, it is enough to check that x_G is contained in the direct sum for any proper flat G of M . If $G \setminus i$ is a not flat of M , then $x_G = \theta_i(x_{G \setminus i})$. If $G \setminus i$ is a flat of M , then either $G \setminus i \in \mathcal{S}_i$ or $G \in \mathcal{S}_i$. In the first case, x_G is an element of the summand indexed by $G \setminus i$. In the second case, $x_G = \theta_i(x_G) - x_{G \cup i} \in \text{CH}_{(i)} + x_{G \cup i} \text{CH}_{(i)}$.

Since our direct sum is a sum of $\text{CH}(M \setminus i)$ -modules and it includes the degree 0 and 1 parts of $\text{CH}(M)$, it will suffice to show that $\text{CH}(M)$ is generated in degrees 0 and 1 as a graded $\text{CH}(M \setminus i)$ -module. In other words, we need to show that

$$\text{CH}_{(i)}^1 \cdot \text{CH}^k(M) = \text{CH}^{k+1}(M) \text{ for any } k \geq 1.$$

We first prove the equality when $k = 1$. Since we have proved that the decomposition (D_1) holds in degree 1, we know that

$$\text{CH}^2(M) = \text{CH}^1(M) \cdot \text{CH}^1(M) = \left(\text{CH}_{(i)}^1 \oplus \bigoplus_{F \in \mathcal{S}_i} \mathbb{Q} x_{F \cup i} \right) \cdot \left(\text{CH}_{(i)}^1 \oplus \bigoplus_{F \in \mathcal{S}_i} \mathbb{Q} x_{F \cup i} \right).$$

Using Lemma 3.8, we may reduce the problem to showing that

$$x_{F \cup i}^2 \in \text{CH}_{(i)}^1 \cdot \text{CH}^1(M) \text{ for any } F \in \mathcal{S}_i.$$

We can rewrite the relation $0 = x_F y_i$ in the augmented Chow ring of M as

$$\begin{aligned}
0 &= (\theta_i(x_F) - x_{F \cup i}) \sum_{i \notin G} x_G \\
&= \theta_i(x_F) \left(\sum_{i \notin G} x_G \right) - x_{F \cup i} \left(\sum_{G \leq F} x_G \right), \\
&= \theta_i(x_F) \left(\sum_{i \notin G} x_G \right) - (\theta_i(x_F) - x_F) \left(\sum_{G < F} x_G \right) - x_{F \cup i} x_F \\
&= \theta_i(x_F) \left(\sum_{i \notin G} x_G - \sum_{G < F} x_G \right) + x_F \left(\sum_{G < F} x_G \right) - x_{F \cup i} \theta_i(x_F) + x_{F \cup i}^2,
\end{aligned}$$

thus reducing the problem to showing that

$$x_F x_G \in \text{CH}_{(i)}^1 \cdot \text{CH}^1(M) \text{ for any } G < F \in \mathcal{S}_i.$$

The collection \mathcal{S}_i is downward closed, meaning that if $G < F \in \mathcal{S}_i$, then $G \in \mathcal{S}_i$; therefore,

$$x_F x_G = (\theta_i(x_F) - x_{F \cup i})(\theta_i(x_G) - x_{G \cup i}).$$

Lemma 3.8 tells us that $x_{F \cup i} x_{G \cup i} \in \text{CH}_{(i)}^1 \cdot \text{CH}^1(M)$, thus so is $x_F x_G$.

We next prove the equality when $k \geq 2$. In this case, we use the result for $k = 1$ along with the fact that the algebra $\text{CH}(M)$ is generated in degree 1 to conclude that

$$\text{CH}_{(i)}^1 \cdot \text{CH}^k(M) = \text{CH}_{(i)}^1 \cdot \text{CH}^1(M) \cdot \text{CH}^{k-1}(M) = \text{CH}^2(M) \cdot \text{CH}^{k-1}(M) = \text{CH}^{k+1}(M).$$

This completes the proof of the decomposition (D_1) and the Poincaré duality theorem for $\text{CH}(M)$ when there is an element i that is not a coloop of M .

The proof when i is a coloop is almost the same; we explain the places where something different must be said. The orthogonality of $x_{E \setminus i} \text{CH}_{(i)}$ and $x_{F \cup i} \text{CH}_{(i)}$ for $F \in \mathcal{S}_i$ follows because $E \setminus i$ and $F \cup i$ are incomparable. To show that the right-hand side of (D_2) spans $\text{CH}(M)$, one extra statement we need to check is that

$$x_{E \setminus i}^2 \in \text{CH}_{(i)}^1 \cdot \text{CH}^1(M).$$

Since i is a coloop, \mathcal{S}_i is the set of all flats properly contained in $E \setminus i$, and we have

$$0 = x_{E \setminus i} y_i = \sum_{i \notin F} x_F x_{E \setminus i} = x_{E \setminus i}^2 + \sum_{F \in \mathcal{S}_i} x_{E \setminus i} x_F = x_{E \setminus i}^2 + \sum_{F \in \mathcal{S}_i} x_{E \setminus i} \theta_i(x_F),$$

where the last equality follows because $E \setminus i$ and $F \cup i$ are not comparable. Thus

$$x_{E \setminus i}^2 = - \sum_{F \in \mathcal{S}_i} x_{E \setminus i} \theta_i(x_F) \in \text{CH}_{(i)}^1 \cdot \text{CH}^1(M).$$

By the induction hypothesis, we know $\text{CH}(M \setminus i)$ satisfies the Poincaré duality theorem. By the coloop case of Lemma 3.3, the Poincaré pairing on $\text{CH}(M)$ restricts to a perfect pairing between

$\text{CH}_{(i)}$ and $x_{E \setminus i} \text{CH}_{(i)}$. Since $\text{CH}_{(i)}$ is a subring of $\text{CH}(M)$ and is zero in degree d , the restriction of the Poincaré pairing on $\text{CH}(M)$ to $\text{CH}_{(i)}$ is zero. Therefore, the subspaces $\text{CH}_{(i)}$ and $x_{E \setminus i} \text{CH}_{(i)}$ intersect trivially, and the restriction of the Poincaré pairing on $\text{CH}(M)$ to $\text{CH}_{(i)} \oplus x_{E \setminus i} \text{CH}_{(i)}$ is non-degenerate. This completes the proof of the theorems about $\text{CH}(M)$ when i is a coloop.

We observe that the surjectivity of the pullback φ_M^\emptyset gives the equality

$$\underline{\text{CH}}_{(i)}^1 \cdot \underline{\text{CH}}^k(M) = \underline{\text{CH}}^{k+1}(M) \text{ for any } k \geq 1.$$

The proof of the theorems about $\underline{\text{CH}}(M)$ then follows by an argument identical to the one used for $\text{CH}(M)$. \square

4. PROOFS OF THE HARD LEFSCHETZ THEOREMS AND THE HODGE–RIEMANN RELATIONS

In this section, we prove Theorem 1.3. Parts (1) and (4) have already been proved in the previous section. We will first prove parts (2) and (3) by induction on the cardinality of E . The proof of parts (5) and (6) is nearly identical to the proof of parts (2) and (3), with the added nuance that we use parts (2) and (3) for the matroid M in the proof of parts (5) and (6) for the matroid M .

For any fan Σ , we will say that Σ satisfies the hard Lefschetz theorem or the Hodge–Riemann relations with respect to some piecewise linear function on Σ if the ring $\text{CH}(\Sigma)$ satisfies the hard Lefschetz theorem or the Hodge–Riemann relations with respect to the corresponding element of $\text{CH}^1(\Sigma)$.

Proof of Theorem 1.3, parts (2) and (3). The statements are trivial when the cardinality of E is 0 or 1, so we will assume throughout the proof that the cardinality of E is at least 2.

Let B be the Boolean matroid on E . By the induction hypothesis, we know that for every nonempty proper flat F of M , the fans $\underline{\Pi}_{M_F}$ and $\underline{\Pi}_{M^F}$ satisfy the hard Lefschetz theorem and the Hodge–Riemann relations with respect to any strictly convex piecewise linear functions on $\underline{\Pi}_{B_F}$ and $\underline{\Pi}_{B^F}$, respectively. By [AHK18, Proposition 7.7], this implies that for every nonempty proper flat F of M , the product $\underline{\Pi}_{M_F} \times \underline{\Pi}_{M^F}$ satisfies the hard Lefschetz theorem and the Hodge–Riemann relations with respect to any strictly convex piecewise linear function on $\underline{\Pi}_{B_F} \times \underline{\Pi}_{B^F}$. In other words, $\underline{\Pi}_M$ satisfies the *local Hodge–Riemann relations* [AHK18, Definition 7.14]:

The star of any ray in $\underline{\Pi}_M$ satisfies the Hodge–Riemann relations.

This in turn implies that $\underline{\Pi}_M$ satisfies the hard Lefschetz theorem with respect to any strictly convex piecewise linear function on $\underline{\Pi}_B$ [AHK18, Proposition 7.15]. It remains to prove only that $\underline{\Pi}_M$ satisfies the Hodge–Riemann relations with respect to any strictly convex piecewise linear function on $\underline{\Pi}_B$.

Let ℓ be a piecewise linear function on $\underline{\Pi}_B$, and let $\underline{\text{HR}}_\ell^k(M)$ be the *Hodge–Riemann form*

$$\underline{\text{HR}}_\ell^k(M) : \underline{\text{CH}}^k(M) \times \underline{\text{CH}}^k(M) \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto (-1)^k \underline{\text{deg}}_M(\ell^{d-2k-1} \eta_1 \eta_2).$$

By [AHK18, Proposition 7.6], the fan $\underline{\Pi}_M$ satisfies the Hodge–Riemann relations with respect to ℓ if and only if, for all $k < \frac{d}{2}$, the Hodge–Riemann form $\underline{\text{HR}}_\ell^k(M)$ is non-degenerate and has the signature

$$\sum_{j=0}^k (-1)^{k-j} \left(\dim \underline{\text{CH}}^j(M) - \dim \underline{\text{CH}}^{j-1}(M) \right).$$

Since $\underline{\Pi}_M$ satisfies the hard Lefschetz theorem with respect to any strictly convex piecewise linear function on $\underline{\Pi}_B$ and signature is a locally constant function on the space of nonsingular forms, the following statements are equivalent:

- (i) The fan $\underline{\Pi}_M$ satisfies the Hodge–Riemann relations with respect to any strictly convex piecewise linear function on $\underline{\Pi}_B$.
- (ii) The fan $\underline{\Pi}_M$ satisfies the Hodge–Riemann relations with respect to some strictly convex piecewise linear function on $\underline{\Pi}_B$.

Furthermore, since satisfying the Hodge–Riemann relations with respect to a given piecewise linear function is an open condition on the function, statement (ii) is equivalent to the following:

- (iii) The fan $\underline{\Pi}_M$ satisfies the Hodge–Riemann relations with respect to some convex piecewise linear function on $\underline{\Pi}_B$.

We show that statement (iii) holds using the semi-small decomposition in Theorem 1.1.

If M is the Boolean matroid B , then $\underline{\text{CH}}(M)$ can be identified with the cohomology ring of the smooth complex projective toric variety $X_{\underline{\Pi}_B}$. Therefore, in this case, Theorem 1.3 is a special case of the usual hard Lefschetz theorem and the Hodge–Riemann relations for smooth complex projective varieties.¹⁶

If M is not the Boolean matroid B , choose an element i that is not a coloop in M , and consider the morphism of fans

$$\pi_i : \underline{\Pi}_M \longrightarrow \underline{\Pi}_{M \setminus i}.$$

By induction, we know that $\underline{\Pi}_{M \setminus i}$ satisfies the Hodge–Riemann relations with respect to any strictly convex piecewise linear function ℓ on $\underline{\Pi}_{B \setminus i}$. We will show that $\underline{\Pi}_M$ satisfies the Hodge–Riemann relations with respect to the pullback $\ell_i := \ell \circ \pi_i$, which is a piecewise linear function on $\underline{\Pi}_B$ that is convex but not necessarily strictly convex.

¹⁶It is not difficult to directly prove the hard Lefschetz theorem and the Hodge–Riemann relations for $\underline{\text{CH}}(B)$ using the coloop case of Theorem 1.1. Alternatively, we may apply McMullen’s hard Lefschetz theorem and Hodge–Riemann relations for polytope algebras [McM93] to the standard permutohedron in \mathbb{R}^E .

By Theorem 1.1, we have the orthogonal decomposition of $\underline{\mathrm{CH}}(M)$ into $\underline{\mathrm{CH}}(M \setminus i)$ -modules

$$\underline{\mathrm{CH}}(M) = \underline{\mathrm{CH}}_{(i)} \oplus \bigoplus_{F \in \mathcal{S}_i} x_{F \cup i} \underline{\mathrm{CH}}_{(i)}.$$

By orthogonality, it is enough to show that each summand of $\underline{\mathrm{CH}}(M)$ satisfies the Hodge–Riemann relations with respect to ℓ_i :

(iv) For every nonnegative integer $k < \frac{d}{2}$, the bilinear form

$$\underline{\mathrm{CH}}_{(i)}^k \times \underline{\mathrm{CH}}_{(i)}^k \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto (-1)^k \underline{\mathrm{deg}}_M(\ell_i^{d-2k-1} \eta_1 \eta_2)$$

is positive definite on the kernel of multiplication by ℓ_i^{d-2k} .

(v) For every nonnegative integer $k < \frac{d}{2}$, the bilinear form

$$x_{F \cup i} \underline{\mathrm{CH}}_{(i)}^{k-1} \times x_{F \cup i} \underline{\mathrm{CH}}_{(i)}^{k-1} \longrightarrow \mathbb{Q}, \quad (\eta_1, \eta_2) \longmapsto (-1)^k \underline{\mathrm{deg}}_M(\ell_i^{d-2k-1} \eta_1 \eta_2)$$

is positive definite on the kernel of multiplication by ℓ_i^{d-2k} .

By Proposition 3.4, the homomorphism θ_i restricts to an isomorphism of $\underline{\mathrm{CH}}(M \setminus i)$ -modules

$$\underline{\mathrm{CH}}(M \setminus i) \cong \underline{\mathrm{CH}}_{(i)}.$$

Thus, statement (iv) follows from Lemma 3.3 and the induction hypothesis applied to $M \setminus i$. By Propositions 2.22, 3.4, and 3.5, the homomorphisms $\theta_i^{F \cup i}$ and $\psi_M^{F \cup i}$ give a $\underline{\mathrm{CH}}(M \setminus i)$ -module isomorphism

$$\underline{\mathrm{CH}}(M_{F \cup i}) \otimes \underline{\mathrm{CH}}(M^F) \cong \underline{\mathrm{CH}}(M_{F \cup i}) \otimes \theta_i^{F \cup i} \underline{\mathrm{CH}}(M^F) \cong x_{F \cup i} \underline{\mathrm{CH}}_{(i)}[1].$$

Note that the pullback of a strictly convex piecewise linear function on $\underline{\Pi}_{B \setminus i}$ to the star

$$\underline{\Pi}_{(B \setminus i)_F} \times \underline{\Pi}_{(B \setminus i)^F} = \underline{\Pi}_{B_{F \cup i}} \times \underline{\Pi}_{B^F}$$

is the class of a strictly convex piecewise linear function. Therefore, statement (v) follows from Proposition 3.7 and the induction applied to $M_{F \cup i}$ and M^F . \square

Proof of Theorem 1.3, parts (5) and (6). This proof is nearly identical to the proof of parts (2) and (3). In that argument, we used the fact that rays of $\underline{\Pi}_M$ are indexed by nonempty proper flats of M and the star of the ray ρ_F is isomorphic to $\underline{\Pi}_{M_F} \times \underline{\Pi}_{M^F}$, which we can show satisfies the hard Lefschetz theorem and the Hodge–Riemann relations using the induction hypothesis. When dealing instead with the augmented Bergman fan $\underline{\Pi}_M$, we have rays indexed by elements of E and rays indexed by proper flats of M , with

$$\mathrm{star}_{\rho_i} \underline{\Pi}_M \cong \underline{\Pi}_{M_{\mathrm{d}(i)}} \quad \text{and} \quad \mathrm{star}_{\rho_F} \underline{\Pi}_M \cong \underline{\Pi}_{M_F} \times \underline{\Pi}_{M^F}.$$

Thus the stars of ρ_i and ρ_F for nonempty F can be shown to satisfy the hard Lefschetz theorem and the Hodge–Riemann relations using the induction hypothesis. However, the star of ρ_\emptyset is isomorphic to $\underline{\Pi}_M$, so we need to use parts (2) and (3) of Theorem 1.3 for M itself. \square

Remark 4.1. It is possible to deduce Poincaré duality, the hard Lefschetz theorem, and the Hodge–Riemann relations for $\text{CH}(M)$ using [AHK18, Theorem 6.19 and Theorem 8.8], where the three properties are proved for generalized Bergman fans $\Sigma_{N,\mathcal{P}}$ in [AHK18, Definition 3.2]. We sketch the argument here, leaving details to the interested readers. Consider the direct sum $M \oplus 0$ of M and the rank 1 matroid on the singleton $\{0\}$ and the order filter $\mathcal{P}(M)$ of all proper flats of $M \oplus 0$ that contain 0. The symbols $B \oplus 0$ and $\mathcal{P}(B)$ are defined in the same way for the Boolean matroid B on E . It is straightforward to check that the linear isomorphism

$$\mathbb{R}^E \longrightarrow \mathbb{R}^{E \cup 0} / \langle \mathbf{e}_E + \mathbf{e}_0 \rangle, \quad \mathbf{e}_j \longmapsto \mathbf{e}_j$$

identifies the complete fan Π_B with the complete fan $\Sigma_{B \oplus 0, \mathcal{P}(B)}$, and the augmented Bergman fan Π_M with a subfan of $\Sigma_{M \oplus 0, \mathcal{P}(M)}$. The third identity in Lemma 2.9 shows that the inclusion of the augmented Bergman fan Π_M into the generalized Bergman fan $\Sigma_{M \oplus 0, \mathcal{P}(M)}$ induces an isomorphism between their Chow rings.

5. PROOF OF THEOREM 1.4

In this section, we prove the decomposition (D₃) by induction on the cardinality of E . The decomposition (D₃) can be proved using the same argument. The results are trivial when E has at most one element. Thus, we may assume that i is an element of E , that $E \setminus i$ is nonempty, and that all the results hold for loopless matroids whose ground set is a proper subset of E .

We first prove that the summands appearing in the right-hand side of (D₃) are orthogonal to each other.

Lemma 5.1. Let F and G be distinct nonempty proper flats of M .

- (1) The spaces $\underline{\psi}_M^F \underline{\text{CH}}(M_F) \otimes \underline{J}_\alpha(M^F)$ and $\underline{H}_\alpha(M)$ are orthogonal in $\underline{\text{CH}}(M)$.
- (2) The spaces $\underline{\psi}_M^F \underline{\text{CH}}(M_F) \otimes \underline{J}_\alpha(M^F)$ and $\underline{\psi}_M^G \underline{\text{CH}}(M_G) \otimes \underline{J}_\alpha(M^G)$ are orthogonal in $\underline{\text{CH}}(M)$.

Proof. The fifth bullet point in Proposition 2.20, together with the fact that $\underline{\psi}_M^F$ is a $\underline{\text{CH}}(M)$ -module homomorphism via $\underline{\varphi}_M^F$, implies that every summand in the right-hand side of (D₃) is an $\underline{H}_\alpha(M)$ -submodule. Thus the first orthogonality follows from the vanishing of $\underline{\psi}_M^F \underline{\text{CH}}(M_F) \otimes \underline{J}_\alpha(M^F)$ in degree $d - 1$.

For the second orthogonality, we may suppose that F is a proper subset of G . Since $\underline{\psi}_M^G$ is a $\underline{\text{CH}}(M)$ -module homomorphism commuting with the degree maps, it is enough to show that

$$\underline{\varphi}_M^G \underline{\psi}_M^F \underline{\text{CH}}(M_F) \otimes \underline{J}_\alpha(M^F) \text{ and } \underline{\text{CH}}(M_G) \otimes \underline{J}_\alpha(M^G) \text{ are orthogonal in } \underline{\text{CH}}(M_G) \otimes \underline{\text{CH}}(M^G).$$

For this, we use the commutative diagram of pullback and pushforward maps

$$\begin{array}{ccc} \underline{\mathrm{CH}}(M_F) \otimes \underline{\mathrm{CH}}(M^F) & \xrightarrow{\psi_M^F} & \underline{\mathrm{CH}}(M) \\ \downarrow \varphi_{M_F}^{G,F} \otimes 1 & & \downarrow \varphi_M^G \\ \underline{\mathrm{CH}}(M_G) \otimes \underline{\mathrm{CH}}(M_F^G) \otimes \underline{\mathrm{CH}}(M^F) & \xrightarrow{1 \otimes \psi_{M^G}^F} & \underline{\mathrm{CH}}(M_G) \otimes \underline{\mathrm{CH}}(M^G), \end{array}$$

which further reduces to the assertion that

$$\psi_{M^G}^F \underline{\mathrm{CH}}(M_F^G) \otimes \underline{J}_\alpha(M^F) \text{ and } \underline{J}_\alpha(M^G) \text{ are orthogonal in } \underline{\mathrm{CH}}(M^G).$$

Since $\underline{J}_\alpha(M^G) \subseteq \underline{H}_\alpha(M^G)$, the above follows from the first orthogonality for M^G . \square

We next show that the restriction of the Poincaré pairing of $\underline{\mathrm{CH}}(M)$ to each summand appearing in the right-hand side of (\underline{D}_3) is non-degenerate.

Lemma 5.2. Let F be a nonempty proper flat of M , and let $k = \mathrm{rk}_M(F)$.

- (1) The restriction of the Poincaré pairing of $\underline{\mathrm{CH}}(M)$ to $\underline{H}_\alpha(M)$ is non-degenerate.
- (2) The restriction of the Poincaré pairing of $\underline{\mathrm{CH}}(M)$ to $\psi_M^F \underline{\mathrm{CH}}(M_F) \otimes \underline{J}_\alpha(M^F)$ is non-degenerate.

Proof. The first statement follows from Proposition 2.26. We prove the second statement.

Since the Poincaré pairing of $\underline{\mathrm{CH}}(M_F)$ is non-degenerate, it is enough to show that the restriction of the Poincaré pairing satisfies

$$\underline{\mathrm{deg}}_M(\psi_M^F(\mu_1 \otimes \nu_1) \cdot \psi_M^F(\mu_2 \otimes \nu_2)) = -\underline{\mathrm{deg}}_{M_F}(\mu_1 \mu_2) \underline{\mathrm{deg}}_{M_F}(\alpha_{M^F} \nu_1 \nu_2).$$

The proof of the identity is nearly identical to that of Proposition 3.7. The left-hand side is

$$\underline{\mathrm{deg}}_M(\psi_M^F(\varphi_{M^F}^F \psi_{M^F}^F(\mu_1 \otimes \nu_1) \cdot (\mu_2 \otimes \nu_2))) = \underline{\mathrm{deg}}_{M_F} \otimes \underline{\mathrm{deg}}_{M^F}(\varphi_{M^F}^F \psi_{M^F}^F(\mu_1 \otimes \nu_1) \cdot (\mu_2 \otimes \nu_2))$$

because ψ_M^F is a $\underline{\mathrm{CH}}(M)$ -module homomorphism commuting with the degree maps. Since the composition $\varphi_{M^F}^F \psi_{M^F}^F$ is multiplication by $\varphi_{M^F}^F(x_F)$, the above becomes

$$-\underline{\mathrm{deg}}_{M_F} \otimes \underline{\mathrm{deg}}_{M^F}((1 \otimes \alpha_{M^F} + \beta_{M^F} \otimes 1) \cdot (\mu_1 \otimes \nu_1) \cdot (\mu_2 \otimes \nu_2)).$$

The vanishing of $\underline{J}_\alpha(M^F)$ in degree $k - 1$ further simplifies the above to the desired expression

$$-\underline{\mathrm{deg}}_{M_F} \otimes \underline{\mathrm{deg}}_{M^F}((1 \otimes \alpha_{M^F}) \cdot (\mu_1 \otimes \nu_1) \cdot (\mu_2 \otimes \nu_2)) = -\underline{\mathrm{deg}}_{M_F}(\mu_1 \mu_2) \underline{\mathrm{deg}}_{M^F}(\alpha_{M^F} \nu_1 \nu_2). \quad \square$$

To complete the proof, we only need to show that the graded vector spaces on both sides of (\underline{D}_3) have the same dimension, which is the next proposition.

Proposition 5.3. As graded vector spaces, there exists an isomorphism

$$\underline{\text{CH}}(\underline{M}) \cong \underline{\text{H}}_{\alpha}(\underline{M}) \oplus \bigoplus_{F \in \underline{\mathcal{C}}(\underline{M})} \underline{\text{CH}}(\underline{M}_F) \otimes \underline{\text{J}}_{\alpha}(\underline{M}^F)[-1], \quad (\underline{\text{D}}'_3)$$

where the sum is over the set $\underline{\mathcal{C}}(\underline{M})$ of proper flats of \underline{M} with rank at least two.

Proof. We prove the proposition using induction on the cardinality of E . Suppose the proposition holds for any matroid whose ground set is a proper subset of E . Suppose that there exists an element $i \in E$ that is not a coloop. Then the decomposition $(\underline{\text{D}}_1)$ implies

$$\underline{\text{CH}}(\underline{M}) \cong \underline{\text{CH}}(\underline{M} \setminus i) \oplus \bigoplus_{G \in \underline{\mathcal{S}}_i(\underline{M})} \underline{\text{CH}}(\underline{M}_{G \cup i}) \otimes \underline{\text{CH}}(\underline{M}^G)[-1],$$

since the maps θ_i , $\theta_i^{G \cup i}$, and $\psi_M^{G \cup i}$ are injective via the Poincaré duality part of Theorem 1.3. By applying the induction hypothesis to the matroids $\underline{M} \setminus i$ and \underline{M}^G , we see that the left-hand side of $(\underline{\text{D}}'_3)$ is isomorphic to the graded vector space

$$\begin{aligned} & \underline{\text{H}}_{\alpha}(\underline{M} \setminus i) \oplus \bigoplus_{G \in \underline{\mathcal{C}}(\underline{M} \setminus i)} \underline{\text{CH}}((\underline{M} \setminus i)_G) \otimes \underline{\text{J}}_{\alpha}((\underline{M} \setminus i)^G)[-1] \\ & \oplus \bigoplus_{G \in \underline{\mathcal{S}}_i(\underline{M})} \bigoplus_{F \in \underline{\mathcal{C}}(\underline{M}^G)} \underline{\text{CH}}(\underline{M}_{G \cup i}) \otimes \underline{\text{H}}_{\alpha}(\underline{M}^G)[-1] \\ & \oplus \bigoplus_{G \in \underline{\mathcal{S}}_i(\underline{M})} \bigoplus_{F \in \underline{\mathcal{C}}(\underline{M}^G)} \underline{\text{CH}}(\underline{M}_{G \cup i}) \otimes \underline{\text{CH}}(\underline{M}_F^G) \otimes \underline{\text{J}}_{\alpha}(\underline{M}^F)[-2]. \end{aligned}$$

Since i is not a coloop, we may replace $\underline{\text{H}}_{\alpha}(\underline{M} \setminus i)$ by $\underline{\text{H}}_{\alpha}(\underline{M})$.

Now, we further decompose the right-hand side of $(\underline{\text{D}}'_3)$ to match the displayed expression. For this, we split the index set $\underline{\mathcal{C}}(\underline{M})$ into three groups:

- (1) $F \in \underline{\mathcal{C}}(\underline{M})$, $i \in F$, $F \setminus i \in \underline{\mathcal{S}}_i(\underline{M})$,
- (2) $F \in \underline{\mathcal{C}}(\underline{M})$, $i \in F$, $F \setminus i \notin \underline{\mathcal{S}}_i(\underline{M})$, and
- (3) $F \in \underline{\mathcal{C}}(\underline{M})$, $i \notin F$.

Suppose F belongs to the first group. In this case, we have $\underline{\text{J}}_{\alpha}(\underline{M}^F) \cong \underline{\text{H}}_{\alpha}(\underline{M}^{F \setminus i})$ as graded vector spaces. Therefore, we have

$$\bigoplus_{\substack{F \in \underline{\mathcal{C}}(\underline{M}) \\ i \in F, F \setminus i \in \underline{\mathcal{S}}_i(\underline{M})}} \underline{\text{CH}}(\underline{M}_F) \otimes \underline{\text{J}}_{\alpha}(\underline{M}^F)[-1] \cong \bigoplus_{G \in \underline{\mathcal{S}}_i(\underline{M})} \underline{\text{CH}}(\underline{M}_{G \cup i}) \otimes \underline{\text{H}}_{\alpha}(\underline{M}^G)[-1].$$

Suppose F belongs to the second group. In this case, $\underline{M}_F = (\underline{M} \setminus i)_{F \setminus i}$ and the matroids \underline{M}^F and $(\underline{M} \setminus i)^{F \setminus i}$ have the same rank. Therefore, we have

$$\bigoplus_{\substack{F \in \underline{\mathcal{C}}(\underline{M}) \\ i \in F, F \setminus i \notin \underline{\mathcal{S}}_i(\underline{M})}} \underline{\text{CH}}(\underline{M}_F) \otimes \underline{\text{J}}_{\alpha}(\underline{M}^F)[-1] \cong \bigoplus_{G \in \underline{\mathcal{C}}(\underline{M} \setminus i) \setminus \underline{\mathcal{C}}(\underline{M})} \underline{\text{CH}}((\underline{M} \setminus i)_G) \otimes \underline{\text{J}}_{\alpha}((\underline{M} \setminus i)^G)[-1].$$

Suppose F belongs to the third group. In this case, we apply (D_1) to M_F and get

$$\begin{aligned}
& \bigoplus_{F \in \underline{\mathcal{C}}(M), i \notin F} \underline{\text{CH}}(M_F) \otimes \underline{\text{J}}_{\alpha}(M^F)[-1] \\
& \cong \bigoplus_{F \in \underline{\mathcal{C}}(M), i \notin F} \left(\underline{\text{CH}}(M_F \setminus i) \oplus \bigoplus_{G \in \underline{\mathcal{S}}_i(M_F)} \underline{\text{CH}}(M_{G \cup i}) \otimes \underline{\text{CH}}(M_F^G)[-1] \right) \otimes \underline{\text{J}}_{\alpha}(M^F)[-1] \\
& \cong \bigoplus_{F \in \underline{\mathcal{C}}(M), i \notin F} \underline{\text{CH}}(M_F \setminus i) \otimes \underline{\text{J}}_{\alpha}(M^F)[-1] \oplus \bigoplus_{\substack{G \in \underline{\mathcal{S}}_i(M) \\ F \in \underline{\mathcal{C}}(M^G)}} \underline{\text{CH}}(M_{G \cup i}) \otimes \underline{\text{CH}}(M_F^G) \otimes \underline{\text{J}}_{\alpha}(M^F)[-2] \\
& \cong \bigoplus_{G \in \underline{\mathcal{C}}(M \setminus i) \cap \underline{\mathcal{C}}(M)} \underline{\text{CH}}((M \setminus i)_G) \otimes \underline{\text{J}}_{\alpha}((M \setminus i)^G)[-1] \oplus \bigoplus_{\substack{G \in \underline{\mathcal{S}}_i(M) \\ F \in \underline{\mathcal{C}}(M^G)}} \underline{\text{CH}}(M_{G \cup i}) \otimes \underline{\text{CH}}(M_F^G) \otimes \underline{\text{J}}_{\alpha}(M^F)[-2].
\end{aligned}$$

The decomposition (D'_3) follows.

Suppose now that every element of E is a coloop of M ; that is, M is a Boolean matroid. We fix an element $i \in E$. The decomposition (D_2) and the Poincaré duality part of Theorem 1.3 imply

$$\underline{\text{CH}}(M) \cong \underline{\text{CH}}(M \setminus i) \oplus \underline{\text{CH}}(M \setminus i)[-1] \oplus \bigoplus_{G \in \underline{\mathcal{S}}_i(M)} \underline{\text{CH}}(M_{G \cup i}) \otimes \underline{\text{CH}}(M^G)[-1].$$

The assumption that i is a coloop implies that $\underline{\mathcal{S}}_i(M) \cap \underline{\mathcal{C}}(M) = \underline{\mathcal{C}}(M \setminus i)$. The induction hypothesis applies to the matroids $M \setminus i$ and M^G , and hence the left-hand side of (D'_3) is isomorphic to

$$\begin{aligned}
& \underline{\text{H}}_{\alpha}(M \setminus i) \oplus \bigoplus_{G \in \underline{\mathcal{C}}(M \setminus i)} \underline{\text{CH}}((M \setminus i)_G) \otimes \underline{\text{J}}_{\alpha}((M \setminus i)^G)[-1] \\
& \oplus \underline{\text{H}}_{\alpha}(M \setminus i)[-1] \oplus \bigoplus_{G \in \underline{\mathcal{C}}(M \setminus i)} \underline{\text{CH}}((M \setminus i)_G) \otimes \underline{\text{J}}_{\alpha}((M \setminus i)^G)[-2] \\
& \oplus \bigoplus_{G \in \underline{\mathcal{S}}_i(M)} \underline{\text{CH}}(M_{G \cup i}) \otimes \left(\underline{\text{H}}_{\alpha}(M^G) \oplus \bigoplus_{F \in \underline{\mathcal{C}}(M^G)} \underline{\text{CH}}(M_F^G) \otimes \underline{\text{J}}_{\alpha}(M^F)[-1] \right)[-1].
\end{aligned}$$

Now, we further decompose the right-hand side of (D'_3) to match the displayed expression. For this, we split the index set $\underline{\mathcal{C}}(M)$ into three groups:

- (1) $F \in \underline{\mathcal{C}}(M)$, $i \in F$,
- (2) $F \in \underline{\mathcal{C}}(M)$, $F = E \setminus i$, and
- (3) $F \in \underline{\mathcal{C}}(M)$, $F \in \underline{\mathcal{S}}_i(M)$.

If F belongs to the first group, then $\underline{\text{J}}_{\alpha}(M^F) \cong \underline{\text{H}}_{\alpha}(M^F \setminus i)$, and hence

$$\bigoplus_{F \in \underline{\mathcal{C}}(M), i \in F} \underline{\text{CH}}(M_F) \otimes \underline{\text{J}}_{\alpha}(M^F)[-1] \cong \bigoplus_{G \in \underline{\mathcal{S}}_i(M)} \underline{\text{CH}}(M_{G \cup i}) \otimes \underline{\text{H}}_{\alpha}(M^G)[-1].$$

If F is the flat $E \setminus i$, we have

$$\underline{\text{H}}_{\alpha}(M) \oplus \underline{\text{CH}}(M_{E \setminus i}) \otimes \underline{\text{J}}_{\alpha}(M^{E \setminus i})[-1] \cong \underline{\text{H}}_{\alpha}(M \setminus i) \oplus \underline{\text{H}}_{\alpha}(M \setminus i)[-1].$$

If F belongs to the third group, we apply (D_2) to M_F and get

$$\begin{aligned}
& \bigoplus_{\substack{F \in \underline{\mathcal{C}}(M) \\ F \in \underline{\mathcal{S}}_i(M)}} \underline{\text{CH}}(M_F) \otimes \underline{J}_\alpha(M^F)[-1] \\
& \cong \bigoplus_{\substack{F \in \underline{\mathcal{C}}(M) \\ F \in \underline{\mathcal{S}}_i(M)}} \left(\underline{\text{CH}}(M_F \setminus i) \oplus \underline{\text{CH}}(M_F \setminus i)[-1] \oplus \bigoplus_{G \in \underline{\mathcal{S}}_i(M_F)} \underline{\text{CH}}(M_{G \cup i}) \otimes \underline{\text{CH}}(M_F^G)[-1] \right) \otimes \underline{J}_\alpha(M^F)[-1] \\
& \cong \bigoplus_{\substack{G \in \underline{\mathcal{C}}(M) \\ G \in \underline{\mathcal{S}}_i(M)}} \underline{\text{CH}}(M_G \setminus i) \otimes \underline{J}_\alpha(M^G)[-1] \oplus \bigoplus_{\substack{G \in \underline{\mathcal{C}}(M) \\ G \in \underline{\mathcal{S}}_i(M)}} \underline{\text{CH}}(M_G \setminus i) \otimes \underline{J}_\alpha(M^G)[-2] \oplus \\
& \quad \bigoplus_{\substack{G \in \underline{\mathcal{S}}_i(M) \\ F \in \underline{\mathcal{C}}(M^G)}} \underline{\text{CH}}(M_{G \cup i}) \otimes \underline{\text{CH}}(M_F^G) \otimes \underline{J}_\alpha(M^F)[-2].
\end{aligned}$$

The decomposition (D'_3) follows. □

Remark 5.4. The decomposition of graded vector spaces appearing in [AHK18, Theorem 6.18] specializes to decompositions of $\underline{\text{CH}}(M)$ and of $\text{CH}(M)$, where the latter goes through Remark 4.1. At the level of Poincaré polynomials, these decompositions coincide with those of Theorem 1.4. However, the subspaces appearing in the decompositions are not the same. In particular, the decompositions in [AHK18, Theorem 6.18] are not orthogonal, and they are not compatible with the $\underline{H}_\alpha(M)$ -module structure on $\underline{\text{CH}}(M)$ or the $H_\alpha(M)$ -module structure on $\text{CH}(M)$.

REFERENCES

- [AHK18] Karim Adiprasito, June Huh, and Eric Katz, *Hodge theory for combinatorial geometries*, Ann. of Math. (2) **188** (2018), no. 2, 381–452. MR3862944 ↑1, 2, 5, 7, 11, 13, 14, 15, 24, 32, 33, 35, 39
- [BDF] Christin Bibby, Graham Denham, and Eva Maria Feichtner, *A Leray model for the Orlik-Solomon algebra*. arXiv:1910.10001. ↑7
- [BES] Spencer Backman, Christopher Eur, and Connor Simpson, *Simplicial generation of Chow rings of matroids*. arXiv:1905.07114. ↑2, 7, 8
- [BHM⁺] Tom Braden, June Huh, Jacob P. Matherne, Nicholas Proudfoot, and Botong Wang, *Singular Hodge theory for combinatorial geometries*. in preparation. ↑2, 4
- [Bjö92] Anders Björner, *The homology and shellability of matroids and geometric lattices*, Matroid applications, 1992, pp. 226–283. MR1165544 ↑9
- [Bri96] Michel Brion, *Piecewise polynomial functions, convex polytopes and enumerative geometry*, Parameter spaces (Warsaw, 1994), 1996, pp. 25–44. MR1481477 ↑14
- [CD06] Michael P. Carr and Satyan L. Devadoss, *Coxeter complexes and graph-associahedra*, Topology Appl. **153** (2006), no. 12, 2155–2168. MR2239078 ↑11
- [CF82] V. P. Camillo and K. R. Fuller, *On graded rings with finiteness conditions*, Proc. Amer. Math. Soc. **86** (1982), no. 1, 1–5. MR663852 ↑29
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. MR2810322 ↑14, 24

- [dCM02] Mark Andrea A. de Cataldo and Luca Migliorini, *The hard Lefschetz theorem and the topology of semismall maps*, Ann. Sci. École Norm. Sup. (4) **35** (2002), no. 5, 759–772. MR1951443 ↑[26](#)
- [dCM09] ———, *The decomposition theorem, perverse sheaves and the topology of algebraic maps*, Bull. Amer. Math. Soc. (N.S.) **46** (2009), no. 4, 535–633. MR2525735 ↑[26](#)
- [Dev09] Satyan L. Devadoss, *A realization of graph associahedra*, Discrete Math. **309** (2009), no. 1, 271–276. MR2479448 ↑[11](#)
- [Ful98] William Fulton, *Intersection theory*, Second, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 2, Springer-Verlag, Berlin, 1998. MR1644323 ↑[14](#)
- [FY04] Eva Maria Feichtner and Sergey Yuzvinsky, *Chow rings of toric varieties defined by atomic lattices*, Invent. Math. **155** (2004), no. 3, 515–536. MR2038195 ↑[2](#), [8](#), [15](#), [17](#)
- [GG82] Robert Gordon and Edward L. Green, *Graded Artin algebras*, J. Algebra **76** (1982), no. 1, 111–137. MR659212 ↑[29](#)
- [Kun86] Joseph P. S. Kung, *Strong maps*, Theory of matroids, 1986, pp. 224–253. MR849397 ↑[10](#)
- [McM93] Peter McMullen, *On simple polytopes*, Invent. Math. **113** (1993), no. 2, 419–444. MR1228132 ↑[33](#)
- [Nel18] Peter Nelson, *Almost all matroids are nonrepresentable*, Bull. Lond. Math. Soc. **50** (2018), no. 2, 245–248. MR3830117 ↑[2](#)
- [Ox11] James Oxley, *Matroid theory*, Second, Oxford Graduate Texts in Mathematics, vol. 21, Oxford University Press, Oxford, 2011. MR2849819 ↑[1](#), [13](#)
- [Pet15] T. Kyle Petersen, *Eulerian numbers*, Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks], Birkhäuser/Springer, New York, 2015. With a foreword by Richard Stanley. MR3408615 ↑[4](#), [16](#)
- [Wel76] D. J. A. Welsh, *Matroid theory*, Academic Press [Harcourt Brace Jovanovich, Publishers], London-New York, 1976. L. M. S. Monographs, No. 8. MR0427112 ↑[1](#)

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS, AMHERST, MA.

E-mail address: `braden@math.umass.edu`

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ.

E-mail address: `junehuh@ias.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR, AND MAX-PLANCK-INSTITUT FÜR MATHEMATIK, BONN, GERMANY.

E-mail address: `matherne@uoregon.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR.

E-mail address: `njp@uoregon.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WI.

E-mail address: `wang@math.wisc.edu`